

CUBIC FORMS, ANOMALY CANCELLATION AND MODULARITY

FEI HAN, RUIZHI HUANG, KEFENG LIU, AND WEIPING ZHANG

ABSTRACT. Motivated by the cubic forms and anomaly cancellation formulas of Witten-Freed-Hopkins, we give some new cubic forms on spin , spin^c , spin^{w_2} and orientable 12-manifolds respectively. We relate them to η -invariants when the manifolds are with boundary, and mod 2 indices on 10 dimensional characteristic submanifolds when the manifolds are spin^c or spin^{w_2} . Our method of producing these cubic forms is a combination of (generalized) Witten classes and the character of the basic representation of affine E_8 .

CONTENTS

Introduction	1
0.1. Spin case	2
0.2. Spin^c and spin^{w_2} cases	5
0.3. Orientable case	6
0.4. Organization of the paper	7
1. Cubic forms on spin 12-manifolds	8
2. Cubic forms on spin^c and spin^{w_2} 12-manifolds	9
2.1. The original cubic forms	9
2.2. New cubic forms on spin^c manifolds	11
3. Cubic forms on orientable 12-manifolds	14
4. Proofs	15
4.1. The basic representation of affine E_8	15
4.2. Proof of Theorem 2.5 and Theorem 2.6	16
4.3. Proof of Theorem 3.1 and Theorem 3.2	22
5. The characteristic classes in the cubic forms	27
6. Obstruction classes of spin^ξ and spin^{w_2} structures	29
References	30

INTRODUCTION

In the recent paper [11], when studying the M-theory anomaly cancellation, Freed and Hopkins developed an algebraic theory of cubic forms. We first briefly review their theory here. Let L be a finitely generated free abelian group. Let

$$(0.1) \quad \langle \cdot, \cdot, \cdot \rangle : L \times L \times L \rightarrow \mathbb{Z}$$

be a symmetric trilinear form on L . For convenience, write the trilinear form simply as a product. $a \in L$ is called a *characteristic element* if the $\bar{a} \in L \otimes \mathbb{Z}/2\mathbb{Z}$ satisfies the following identity

$$(0.2) \quad \bar{a} \bar{x} \bar{y} \equiv \bar{x} \bar{x} \bar{y} + \bar{x} \bar{y} \bar{y} \pmod{2}$$

for any $\bar{x}, \bar{y} \in L \otimes \mathbb{Z}/2\mathbb{Z}$. Let $L_{\text{char}} \subset L$ be the torsor of characteristic elements in L . Let $L^* = \text{Hom}(L; \mathbb{Z})$.

Theorem 0.1 (Freed-Hopkins, Lemma 4.13 in [11]). *Let $a \in L_{\text{char}}$ and \hat{a} be the mod 24 reduction. There exists a unique $\hat{b} \in L^* \otimes \mathbb{Z}/24\mathbb{Z}$ such that*

$$(0.3) \quad \hat{b}(\hat{x}) = 4\hat{x}^3 + 6\hat{a}\hat{x}^2 + 3\hat{a}^2\hat{x} \pmod{24}$$

for all $\hat{x} \in L \otimes \mathbb{Z}/24\mathbb{Z}$.

Let $h(x)$ be a degree 3 polynomial whose highest term is $\frac{1}{6}x^3$. Then the following algebraic identity shows that h is a cubic refinement of the trilinear form xyz :

$$(0.4) \quad xyz = h(x+y+z) - h(x+y) - h(x+z) - h(y+z) + h(x) + h(y) + h(z) - h(0).$$

Let $a \in L, b \in L^*$. Consider the Witten-Freed-Hopkins polynomial on L :

$$(0.5) \quad f_{a,b}(x) = (a+x)^3 - b(a+x).$$

It is easy to see that

$$(0.6) \quad \frac{f_{a,b}(2x) - f_{a,b}(0)}{48}$$

is a degree 3 polynomial whose highest term is $\frac{1}{6}x^3$, and therefore a cubic refinement of xyz ; moreover when a is a characteristic element and b satisfies (0.3),

$$(0.7) \quad \frac{f_{a,b}(2x) - f_{a,b}(0)}{48} \in \mathbb{Z}.$$

0.1. Spin case. Let Z be a 12 dimensional smooth manifold. Denote the integral Pontrjagin classes and the Stiefel-Whitney classes of Z by p_i, w_i respectively. Let $x \in H^4(Z; \mathbb{Z})$. Following Witten [31], x determines an isomorphism class of principal E_8 bundles on Z . Let $V(x)$ denote the real adjoint vector bundle associated to the principal E_8 bundle determined by the class x . Denote by $V_{\mathbb{C}}(x)$ the complexification of $V(x)$. The Chern character of $V_{\mathbb{C}}(x)$ is (c.f. (4.31) in [11])

$$(0.8) \quad \text{ch}(V_{\mathbb{C}}(x)) = 248 - 60x + 6x^2 - \frac{1}{3}x^3.$$

So by the expression of the Chern character in terms of the Chern classes, it is easy to see that

$$(0.9) \quad x = \frac{1}{60}c_2(V_{\mathbb{C}}(x)).$$

Suppose Z is closed and oriented. Let $L = H^4(Z; \mathbb{Z})$ and the trilinear form is the cup product of three elements in L evaluated on the fundamental class $[Z]$.

Further suppose Z is spin. There is a canonical degree 4 class $\lambda \in H^4(Z; \mathbb{Z})$ such that $2\lambda = p_1$ and $\lambda \equiv w_4 \pmod{2}$ ([11]). The following theorem shows that λ is a characteristic element.

Theorem 0.2 (Freed-Hopkins, Lemma 4.22 in [11]). *The Stiefel-Whitney class $\bar{\lambda} = w_4$ of a closed spin 12-manifold satisfies (0.2) and thus λ is a characteristic element.*

For $a = \lambda$, Freed and Hopkins showed that there is a characteristic class $p \in H^8(Z; \mathbb{Z})$ such that $2p = p_2 - \lambda^2$ and when $b = p$, (0.3) is satisfied. This is deduced from the beautiful and important anomaly cancellation formula discovered by Witten-Freed-Hopkins (Theorem 1.2), which is proved by a direct computation, as well as the famous Atiyah-Hirzebruch divisibility on $8k+4$ dimensional spin manifolds. Adopting the above notations, one sees that on a closed spin smooth 12-manifold Z ,

$$(0.10) \quad \frac{f_{\lambda,p}(2x)}{48}$$

is a half integer, which has analytic meaning as (quarter of) indices of certain twisted Dirac operators on Z (Theorem 1.2).

The aim of this paper is to show how the Witten-Freed-Hopkins anomaly cancellation formula as well as several interesting generalizations can be derived in a uniform way from the extended theory of the Witten genus.

In fact, let M be a $4m$ dimensional compact oriented smooth manifold. Let

$$\{\pm 2\pi\sqrt{-1}z_j, 1 \leq j \leq 2m\}$$

denote the formal Chern roots of $T_{\mathbb{C}}M$, the complexification of the tangent vector bundle TM of M . The famous Witten genus of M can be written as

$$W(M) = \left\langle \left(\prod_{j=1}^{2m} z_j \frac{\theta'(0, \tau)}{\theta(z_j, \tau)} \right), [M] \right\rangle \in \mathbb{Q}[[q]],$$

with $\tau \in \mathbb{H}$, the upper half-plane, and $q = e^{2\pi\sqrt{-1}\tau}$. The Witten genus was first introduced in [30] and can be viewed as the loop space analogue of the \hat{A} -genus. It can be expressed as a q -deformed \hat{A} -genus as

$$W(M) = \left\langle \hat{A}(TM) \text{ch}(\Theta(T_{\mathbb{C}}M)), [M] \right\rangle,$$

where

$$\Theta(T_{\mathbb{C}}M) = \bigotimes_{n=1}^{\infty} S_{q^n}(\widetilde{T_{\mathbb{C}}M}), \quad \text{with } \widetilde{T_{\mathbb{C}}M} = T_{\mathbb{C}}M - \mathbb{C}^{4m},$$

is the Witten bundle introduced in [30]. When the manifold M is spin, according to the Atiyah-Singer index theorem, the Witten genus can be expressed analytically as the index of the twisted Dirac operator,

$$W(M) = \text{Ind}(D \otimes \Theta(T_{\mathbb{C}}M)) \in \mathbb{Z}[[q]],$$

where D is the Atiyah-Singer spin Dirac operator on M . Moreover, if M is string, i.e.

$$\lambda = \frac{1}{2}p_1(TM) = 0,$$

or even weaker, if M is spin and the first rational Pontrjagin class of M vanishes, then $W(M)$ is a modular form of weight $2m$ over $SL(2, \mathbb{Z})$ with integral Fourier development ([32]). The homotopy theoretical refinement of the Witten genus on string manifolds leads to the beautiful theory of tmf (*topological modular form*) developed by Hopkins and Miller [21]. The string condition is the orientability condition for this generalized cohomology theory.

If the string condition $\lambda = 0$ does not hold, one constructs the cohomology class (c.f. [15]),

$$(0.11) \quad \mathcal{W}(TM) = e^{\frac{1}{24}E_2(\tau) \cdot p_1(TM)} \hat{A}(TM) \text{ch}(\Theta(T_{\mathbb{C}}M)) \in H^{4*}(M; \mathbb{Q})[[q]],$$

where $E_2(\tau)$ is the Eisenstein series of weight 2 (c.f. Chap 2.3 in [7]). We call $\mathcal{W}(TM)$ the *Witten class* of M

Let P be a principal E_8 bundle over M . In Section 4.2.1, we consider an associated element $\mathcal{V} \in K(M)[[q]]$ constructed from the basic representation of affine E_8 . Let P_1, P_2 be two principal E_8 bundles with the corresponding $\mathcal{V}_1, \mathcal{V}_2$. Let W_1, W_2 be the complexified vector bundles associated to the adjoint representation of E_8 . Denote

$$\varphi(\tau) = \prod_{n=1}^{\infty} (1 - q^n).$$

We construct the *twisted Witten class* (specializing to ξ trivial and $c = 0$ in (4.18) for the spin case here for simplicity)

$$(0.12) \quad \mathcal{Q}(\mathcal{V}_1, \mathcal{V}_2) := e^{\frac{1}{24}E_2(\tau) \cdot (\frac{1}{30}(c_2(W_1) + c_2(W_2)))} \mathcal{W}(TM) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_1) \text{ch}(\mathcal{V}_2) \in H^{4*}(M; \mathbb{Q})[[q]]$$

and show that the degree 12 component is a modular form of weight 14 over $SL(2, \mathbb{Z})$ when $M = Z$ is 12 dimensional. Using the fact that the space of modular forms of weight 14 over $SL(2, \mathbb{Z})$ is

1-dimensional and spanned by $E_4^2(\tau)E_6(\tau)$, where $E_4(\tau)$ and $E_6(\tau)$ are the Eisenstein series of weight 4 and 6 respectively (c.f. Chap 2.1 in [7]), we deduce a factorization formula (4.33). This formula reduces to the Witten-Freed-Hopkins anomaly cancellation formula (1.1) (when ξ is trivial and $c = 0$).

Back to the algebraic theory of cubic forms, consider the polynomial

$$(0.13) \quad \tilde{f}_{a,b}(x) = 4(a+x)^3 - 6a(a+x)^2 - (b-3a^2)(a+x)$$

with understanding $3a^2$ as an element in L^* by abusing notations.

It is easy to check that

$$(0.14) \quad \tilde{f}_{a,b}(x) = \frac{f_{a,b}(2x) + f_{a,b}(0)}{2}.$$

From (0.4), (0.6) and (0.14), we see that $\frac{\tilde{f}_{a,b}(x) - \tilde{f}_{a,b}(0)}{24}$ is a cubic refinement of xyz , and when a is a characteristic element and b modulo 24 satisfies (0.3), the following holds,

$$(0.15) \quad \frac{\tilde{f}_{a,b}(x) - \tilde{f}_{a,b}(0)}{24} \in \mathbb{Z}.$$

In view of (0.14) and the half integrality of $\frac{f_{\lambda,p}(2x)}{48}$ as well as its analytic meaning, we see that on a closed 12 dimensional spin manifold Z , when $a = \lambda, b = p$,

$$(0.16) \quad \frac{\tilde{f}_{\lambda,p}(x)}{24}$$

is an integer, which has analytic meaning as (half of) the indices of certain twisted Dirac operator on Z (Theorem 1.3).

In Section 4.2.2, we show that Theorem 1.3 can actually also be deduced from a factorization formula (4.42) proved there by constructing the *twisted Witten class* (specializing to ξ trivial and $c = 0$ in (4.35) for the spin case here for simplicity)

$$(0.17) \quad \mathcal{R}(\mathcal{V}) = e^{\frac{1}{24}E_2(\tau) \cdot \frac{1}{30}c_2(W)} \mathcal{W}(TZ) \varphi(\tau)^8 \text{ch}(\mathcal{V}) \in H^{4*}(Z; \mathbb{Q})[[q]].$$

To consider $\tilde{f}_{\lambda,p}(x)$ defined via (0.14) might look redundant. However we include it here because first it arises from the modular form $\mathcal{R}(\mathcal{V})$, different from $\mathcal{Q}(\mathcal{V}_1, \mathcal{V}_2)$; secondly, we would like to point out that the relation (0.14) between the cubic form $\frac{\tilde{f}_{\lambda,p}(x)}{24}$ and the cubic form $\frac{f_{\lambda,p}(2x)}{48}$ corresponds exactly to the relation between the corresponding modular forms:

$$(0.18) \quad \mathcal{R}(\mathcal{V}) = \sqrt{\mathcal{Q}(\mathcal{V}, \mathcal{V}) \cdot \mathcal{W}(\overline{TZ})} \in H^{4*}(Z; \mathbb{Q})[[q]].$$

Remark 0.1. *The method of constructing $\mathcal{Q}(\mathcal{V}_1, \mathcal{V}_2)$, $\mathcal{R}(\mathcal{V})$ and using their modularities to prove factorization formulas appeared in [19] for fiber bundles with 10 dimensional fibers. In this paper, we apply this method to 12 dimensional manifolds.*

Remark 0.2. *In [11], Freed and Hopkins also studied anomaly cancellation on pin^+ manifolds. For a pin^+ 12-manifold Z with \mathfrak{m}_c structure (w_1 -twisted integral lift of w_4), they showed that there exists a characteristic class $\tilde{c} \in H^4(Z; \tilde{\mathbb{Z}})/\text{torsion}$, which is a characteristic element in*

$$L = H^4(Z; \tilde{\mathbb{Z}})/\text{torsion}$$

and a characteristic class $\bar{p} \in H^8(Z; \mathbb{Z})/\text{torsion}$ such that (0.3) is satisfied. Then

$$(0.19) \quad \frac{f_{\tilde{c}, \bar{p}}(0)}{48}$$

is a half integer. Freed-Hopkins (Theorem 2.2, Theorem 6.3 in [11]) proved the following anomaly cancellation

$$(0.20) \quad \exp\left(-2\pi i \cdot \frac{f_{\tilde{c}, \tilde{p}}(0)}{48}\right) \cdot \exp\left(2\pi i \cdot \frac{\eta(TZ - 2)}{4}\right) = 1.$$

We have not recovered this result via modularity yet.

0.2. Spin^c and spin^{w2} cases. Suppose Z is a closed 12 dimensional smooth manifold not necessarily spin. Then the classes λ and p in the previous subsection for the spin case do not necessarily exist. By abusing notations we just denote $\frac{1}{2}p_1$ by λ , and $\frac{1}{2}(p_2 - \frac{1}{4}p_1^2)$ by p in $H^*(Z; \mathbb{Q})$. The original cubic forms $\frac{f_{\lambda, p}(2x)}{48}$ and $\frac{\tilde{f}_{\lambda, p}(x)}{24}$ now only take values in rationals rather than integers and lose analytic interpretations as indices of twisted Dirac operators.

In Section 2.1, we will show that the analytic interpretations can be rescued on spin^c and spin^{w2} manifolds by applying the Rokhlin congruence formulas established in [33, 34, 36, 37] via mod 2 indices on 10 dimensional spin or pin⁻ manifolds. Let us be more precise in the following.

Let K be an $8k + 4$ dimensional spin^c manifold. Let ξ be the complex line bundle of the spin^c structure. Let $c = c_1(\xi) \in H^2(K; \mathbb{Z})$. Let U be a *characteristic submanifold* of the spin^c structure, i.e. an orientable $8k + 2$ dimensional submanifold of K such that $[U] \in H_{8k+2}(K; \mathbb{Z})$ is dual to c . U carries a canonically induced spin structure up to spin cobordism. Let D_U be the Atiyah-Singer spin Dirac operator on U .

The Rokhlin congruence formula (2.2) in Theorem 2.1 allows one to write the twisted \hat{A} -genus on K in terms of mod 2 indices of twisted Dirac operators on U with a correction term. Combining (2.2) with the Witten-Freed-Hopkins anomaly cancellation formula (1.1) and the new formula (1.2), we obtain Theorem 2.2.

We call an $8k + 4$ dimensional closed smooth oriented manifold K a *spin^{w2} manifold* if there exists a rank 2 nonorientable real vector bundle ξ such that $w_2(TK) = w_2(\xi)$. For such manifolds, the corresponding Rokhlin congruence formula has been studied in [34]. Let U be a *characteristic submanifold* of the spin^{w2} structure, i.e. a nonorientable $8k + 2$ dimensional submanifold of K such that $[U] \in H_{8k+2}(K; \mathbb{Z}_2)$ is dual to $w_2(TK) \in H^2(K; \mathbb{Z}_2)$. U carries a canonically induced pin⁻ structure up to pin⁻ cobordism.

The Rokhlin congruence formula (2.5) in Theorem 2.3 allows one to write the twisted \hat{A} -genus of K in terms of twisted mod 2 analytic indices on U with a correction term. Combining (2.5) with the Witten-Freed-Hopkins anomaly cancellation formula (1.1) and the new formula (1.2), we obtain Theorem 2.4. Details about spin^{w2} structures and the obstruction classes to them will be studied in Section 6.

Section 2.2 provides another way on spin^c manifolds to restore the beautiful nature of the Witten-Freed-Hopkins anomaly cancellation formula on spin manifolds. Generally there are no characteristic elements on spin^c manifolds like λ for spin manifolds. In the algebraic theory of cubic forms, suppose $a \in L$ is not necessarily a characteristic element, then the Freed-Hopkins Theorem 0.1 is weakened to mod 12. More precisely, if \hat{a} is the mod 12 reduction, then there exists a unique $\hat{b} \in L^* \otimes \mathbb{Z}/12\mathbb{Z}$ such that

$$(0.21) \quad \hat{b}(\hat{x}) = 4\hat{x}^3 + 6\hat{a}\hat{x}^2 + 3\hat{a}^2\hat{x} \pmod{12}$$

for all $\hat{x} \in L \otimes \mathbb{Z}/12\mathbb{Z}$.

Let Z be a 12 dimensional closed smooth spin^c manifold. Let c be the first Chern class of the complex line bundle of the spin^c structure. In Section 2.2, we will show that on Z there exist characteristic classes $a = \lambda_c \in H^4(Z; \mathbb{Z})$, $b = p_c \in H^8(Z; \mathbb{Z})$ such that (0.21) holds. This is derived from Theorem 2.5 and Theorem 2.6, in which we show that

$$(0.22) \quad \frac{f_{\lambda_c, p_c}(2x)}{24}$$

is a half integer and

$$(0.23) \quad \frac{\tilde{f}_{\lambda_c, p_c}(x)}{12}$$

is an integer, by demonstrating their analytic meanings using indices of twisted spin^c Dirac operators.

The left hand sides of Theorem 2.5 and Theorem 2.6 provide new cubic forms on spin^c manifolds, generalizing the cubic forms in the spin case when the manifold is spin and $c = 0$. The coefficients appearing in the new cubic forms will be studied in Section 5.

Clearly

$$h(x) = \frac{f_{\lambda_c, p_c}(2x) - f_{\lambda_c, p_c}(0)}{24}$$

is a polynomial of x valued in \mathbb{Z} with highest term $\frac{1}{3}x^3$. We therefore see that on spin^c 12-manifolds, there is an integral cubic refinement only for $2xyz$ rather than xyz ,

$$(0.24) \quad 2\langle x \cup y \cup z, [Z] \rangle = h(x+y+z) - h(x+y) - h(x+z) - h(y+z) + h(x) + h(y) + h(z) - h(0).$$

Theorem 2.5 and Theorem 2.6 are deduced from the factorization formulas (4.33) and (4.42), which are proved in Section 4.2 by constructing the generalized Witten class

$$(0.25) \quad \mathcal{W}_c(TZ) := e^{\frac{1}{24}E_2(\tau) \cdot (p_1(TZ) - 3c^2)} \hat{A}(TZ) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \in H^{4*}(Z; \mathbb{Q})[[q]]$$

and the *twisted generalized Witten classes*

$$(0.26) \quad \mathcal{Q}_c(\mathcal{V}_1, \mathcal{V}_2) := e^{\frac{1}{24}E_2(\tau) \cdot (\frac{1}{30}(c_2(W_1) + c_2(W_2)))} \mathcal{W}_c(TZ) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_1) \text{ch}(\mathcal{V}_2) \in H^{4*}(Z; \mathbb{Q})[[q]],$$

$$(0.27) \quad \mathcal{R}_c(\mathcal{V}) := e^{\frac{1}{24}E_2(\tau) \cdot \frac{1}{30}c_2(W)} \mathcal{W}_c(TZ) \varphi(\tau)^8 \text{ch}(\mathcal{V}) \in H^{4*}(Z; \mathbb{Q})[[q]].$$

Applying the Rokhlin congruence (2.1) in Theorem 2.1 to Theorems 2.5 and 2.6, we give mod 2 index interpretations of $\frac{f_{\lambda_c, p_c}(2x)}{12}$ and $\frac{\tilde{f}_{\lambda_c, p_c}(x)}{12}$ in Theorem 2.7. Now on spin^c manifolds, we have two types of mod 2 formulas: Theorem 2.2 and Theorem 2.7. Subtracting the corresponding sides of these formulas, we obtain Corollary 2.1, which involves interesting quadratic forms on 10 dimensional spin manifolds and mod 2 indices. See Remark 2.1.

0.3. Orientable case. Let Z be a 12 dimensional closed smooth oriented manifold without assuming any additional topological constraints. In this general situation, we are not able to find characteristic classes a of degree 4 and b of degree 8 such that the mod 12 equality (0.21) holds for all $x \in H^4(Z; \mathbb{Z})$.

However, by our modularity method, we find that if

$$b = 4p_1^2 - 7p_2$$

and $\hat{b} = b \pmod{3}$, then the following mod 3 equality holds,

$$(0.28) \quad \hat{b}(\hat{x}) = 4\hat{x}^3 \pmod{3}$$

for all $\hat{x} \in H^4(Z; \mathbb{Z}) \otimes \mathbb{Z}/3\mathbb{Z}$. In fact, in Theorem 3.1 and Theorem 3.2, we will show that when $a = -p_1$,

$$(0.29) \quad \frac{f_{a,b}(2x)}{6}$$

is a half integer and

$$(0.30) \quad \frac{\tilde{f}_{a,b}(x)}{3}$$

is an integer, by demonstrating their analytic meanings as indices of twisted signature operators.

Now let

$$h(x) = \frac{f_{a,b}(2x) - f_{a,b}(0)}{6},$$

which is a polynomial of x with value in \mathbb{Z} and highest term $\frac{8}{6}x^3$. We therefore see that on general oriented 12-manifolds, there is an integral cubic refinement only for $8xyz$ rather than for $2xyz$ or xyz ,

$$(0.31) \quad 8\langle x \cup y \cup z, [Z] \rangle = h(x+y+z) - h(x+y) - h(x+z) - h(y+z) + h(x) + h(y) + h(z) - h(0).$$

We prove Theorem 3.1 and Theorem 3.2 in Section 4.3 by constructing the \widehat{L} -Witten class and twisted \widehat{L} -Witten classes (see (4.47), (4.49) and (4.61)).

Remark 0.3. *In view of Theorem 1 in [26], we may obtain formulas more general than the ones presented in Section 3. In fact, let F be a spin vector bundle of even rank over M such that*

$$\frac{1}{2}p_1(F) = \frac{1}{2}p_1(M)$$

and $S^\pm(F)$ be the spinor bundles of F . In the construction of the \widehat{L} -Witten class (4.47), one can replace the bundle

$$\Theta_1(T_{\mathbb{C}}M) \otimes \Theta_2(T_{\mathbb{C}}M) \otimes \Theta_3(T_{\mathbb{C}}M)$$

by

$$\Theta_1(F_{\mathbb{C}}) \otimes \Theta_2(F_{\mathbb{C}}) \otimes \Theta_3(F_{\mathbb{C}})$$

(see the construction of $\Theta_1, \Theta_2, \Theta_3$ in (4.43), (4.44), (4.45)). Then the similar modularity method will deduce formulas giving analytic interpretations to some interesting new cubic forms (depending on the rank of F) via the indices of twisted Dirac operators

$$\begin{aligned} & D \otimes ((S^+(F) \oplus S^-(F)) \otimes V_{\mathbb{C}}(x)), \\ & D \otimes ((S^+(F) \oplus S^-(F)) \otimes T_{\mathbb{C}}Z), \\ & D \otimes ((S^+(F) \oplus S^-(F)) \otimes (\wedge^2(F_{\mathbb{C}}) - S^2(F_{\mathbb{C}}))), \end{aligned}$$

and

$$D \otimes (S^+(F) \oplus S^-(F)).$$

0.4. Organization of the paper. In Section 1, we review the Witten-Freed-Hopkins formula (Theorem 1.2) and present the new type of anomaly cancellation formula (Theorem 1.3). We point out that they are special cases of the corresponding anomaly cancellation formulas for spin^c manifolds, which are given in Section 2.2. As these formulas are consequences of the factorization formulas (4.33) and (4.42), which hold on the level of differential forms, Theorems 1.2 and 1.3 have analogues for manifolds with boundary. They are stated in Theorem 1.4.

In Section 2, we give the Witten-Freed-Hopkins anomaly cancellation formulas on 12 dimensional spin^c and spin^{w_2} manifolds. First in Section 2.1, we consider the original cubic forms as in the spin case. We use the Rokhlin congruence formulas in [33, 34, 36, 37] to express the original cubic forms as mod 2 indices on 10 dimensional characteristic submanifolds with correction terms. Then in Section 2.2, for the spin^c case, we give new cubic forms and anomaly cancellation formulas, which generalize the anomaly cancellation formulas in the spin case. We will also give the mod 2 congruence formulas for the new cubic forms as well as the formulas for manifolds with boundary.

In Section 3, we present the anomaly cancellation formulas for 12 dimensional orientable manifolds.

In Section 4, the proofs of the main theorems (Theorems 2.5, 2.6, 3.1 and 3.2) in Section 2 and Section 3 will be given. Actually what we prove are the factorization formulas (4.33), (4.42), (4.59) and (4.69), which are all equalities on the levels of differential forms.

In Section 5, we study the characteristic class coefficients appearing in the cubic forms in Theorems 2.5, 2.6, 3.1 and 3.2.

In Section 6, details about spin^{w_2} structures and the obstruction classes to them are studied.

Acknowledgements. Fei Han was partially supported by the grant AcRF R-146-000-263-114 from National University of Singapore.

Ruizhi Huang was supported by Postdoctoral International Exchange Program for Incoming Postdoctoral Students under Chinese Postdoctoral Council and Chinese Postdoctoral Science Foundation. He was also supported in part by Chinese Postdoctoral Science Foundation (Grant nos. 2018M631605 and 2019T120145), and National Natural Science Foundation of China (Grant no. 11801544).

Kefeng Liu is partially supported by an NSF grant.

Weiping Zhang was partially supported by the NNSFC Grant no. 11931007.

1. CUBIC FORMS ON SPIN 12-MANIFOLDS

In this section, we review the Witten-Freed-Hopkins anomaly cancellation formula, present the new type of cancellation formula and point out that they can be deduced from the more general formulas for the spin^c case in Section 2.2. We will also present the corresponding formulas when the manifolds have boundary.

Let Z be a closed spin smooth 12-manifold. Recall that $\lambda \in H^4(Z; \mathbb{Z})$ satisfies $2\lambda = p_1$.

Theorem 1.1 (Freed-Hopkins [11]). *There is a canonical degree 8 integral class $p \equiv w_8 \pmod{2}$ such that*

$$2p = p_2 - \lambda^2.$$

Let $V(x)$ denote the real adjoint vector bundle associated to the principal E_8 bundle determined by a class $x \in H^4(Z; \mathbb{Z})$. Denote by $V_{\mathbb{C}}(x)$ the complexification of $V(x)$. Let

$$C(x) = \lambda + 2x \in H^4(Z; \mathbb{Z}).$$

One has the following important formula,

Theorem 1.2 (Witten-Freed-Hopkins [31, 11]). *The following identity holds,*

$$(1.1) \quad \left\langle \frac{C(x)[p - C(x)^2]}{48}, [Z] \right\rangle = \left\langle \frac{1}{2} \hat{A}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + \frac{1}{4} \hat{A}(TZ) \text{ch}(T_{\mathbb{C}}Z) - \hat{A}(TZ), [Z] \right\rangle.$$

Proof. Taking $\xi = \mathbb{C}$ and $c = 0$ in Theorem 2.5, one obtains (1.1). □

We also have the following new cancellation formula. Let

$$\tilde{C}(x) = \lambda + x \in H^4(Z; \mathbb{Z}).$$

Theorem 1.3. *Let*

$$\tilde{p} = p - 3\lambda^2.$$

The following identity holds,

$$(1.2) \quad \left\langle \frac{\tilde{C}(x)[\tilde{p} + 6\lambda\tilde{C}(x) - 4\tilde{C}(x)^2]}{24}, [Z] \right\rangle = \left\langle \frac{1}{2} \hat{A}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + \frac{1}{2} \hat{A}(TZ) \text{ch}(T_{\mathbb{C}}Z) + 122 \hat{A}(TZ), [Z] \right\rangle.$$

Proof. Taking $\xi = \mathbb{C}$ and $c = 0$ in Theorem 2.6, one obtains (1.2). □

Now suppose Z has boundary and let Y be the boundary of Z with the induced spin structure. Let $g^{TZ}, g^{V(x)}$ be a Riemannian metric on TZ and a Euclidean metric on $V(x)$ respectively. Let ∇^{TZ} be the Levi-Civita connection on TZ and $\nabla^{V(x)}$ be a Euclidean connection on $V(x)$. $g^{TZ}, g^{V(x)}, \nabla^{TZ}, \nabla^{V(x)}$ induce the corresponding Hermitian metrics and connections on $T_{\mathbb{C}}Z$ and $V_{\mathbb{C}}(x)$, the complexifications. Assume all the involved metrics and connections are of product structures near $\partial Z = Y$.

Let D_Y be the Atiyah-Singer Dirac operator on Y . Let $\bar{\eta}$ denote the reduced η -invariant in the sense of Atiyah-Patodi-Singer [2].

Denote by $p_i(\nabla^{TZ})$ the i -th Pontrjagin form of (TZ, ∇^{TZ}) (c.f. [35]). Denote by $\lambda(\nabla^{TZ})$ the characteristic form $\frac{1}{2}p_1(\nabla^{TZ})$ and by $p(\nabla^{TZ})$ the characteristic form $\frac{1}{2}p_2(\nabla^{TZ}) - \frac{1}{8}p_1(\nabla^{TZ})^2$.

In the following, when a connection appears in a bracket of a characteristic class, we always mean the corresponding characteristic form determined by this connection.

Denote

$$\check{x} = \frac{1}{60}c_2(V_{\mathbb{C}}(x), \nabla^{V_{\mathbb{C}}(x)}).$$

Let

$$C(\check{x}) = \lambda(\nabla^{TZ}) + 2\check{x}$$

and

$$\tilde{C}(\check{x}) = \lambda(\nabla^{TZ}) + \check{x}.$$

As (1.1) and (1.2) hold on the level of forms, by the Atiyah-Patodi-Singer index theorem [2], we have the following formulas.

Theorem 1.4.

$$(1.3) \quad \frac{1}{24} \int_Z C(\check{x})[p(\nabla^{TZ}) - C(\check{x})^2] \equiv \bar{\eta}(D_Y^{V_{\mathbb{C}}(x)}) + \frac{1}{2}\bar{\eta}(D_Y^{T_{\mathbb{C}}Z}) - 2\bar{\eta}(D_Y) \pmod{\mathbb{Z}};$$

and

$$(1.4) \quad \frac{1}{24} \int_Z \tilde{C}(\check{x})[\tilde{p}(\nabla^{TZ}) + 6\lambda(\nabla^{TZ})\tilde{C}(\check{x}) - 4\tilde{C}(\check{x})^2] \equiv \frac{1}{2}\bar{\eta}(D_Y^{V_{\mathbb{C}}(x)}) + \frac{1}{2}\bar{\eta}(D_Y^{T_{\mathbb{C}}Z}) + 122\bar{\eta}(D_Y) \pmod{\mathbb{Z}}.$$

2. CUBIC FORMS ON spin^c AND spin^{w_2} 12-MANIFOLDS

In this section, we extend the Witten-Freed-Hopkins anomaly cancellation formulas to 12 dimensional spin^c and spin^{w_2} manifolds.

2.1. The original cubic forms. Suppose Z is a closed 12 dimensional smooth manifold not necessarily spin. Then the characteristic classes λ and p in the above section for the spin case do not necessarily exist. By abusing notations we simply denote $\frac{1}{2}p_1$ by λ , and $\frac{1}{2}(p_2 - \frac{1}{4}p_1^2)$ by p in $H^*(Z; \mathbb{Q})$.

Let $x \in H^4(Z; \mathbb{Z})$. Let $V(x)$ and $V_{\mathbb{C}}(x)$ be the same meaning as introduced in the beginning of Section 0.1.

2.1.1. spin^c case. Let K be an $8k+4$ dimensional spin^c manifold. Let ξ be the complex line bundle of the spin^c structure. Let $c = c_1(\xi) \in H^2(K; \mathbb{Z})$. Let U be a *characteristic submanifold* of the spin^c structure, i.e. an orientable $8k+2$ dimensional submanifold of K such that $[U] \in H_{8k+2}(K; \mathbb{Z})$ is dual to c . U carries a canonically induced spin structure up to spin cobordism. Let D_U be the Atiyah-Singer spin Dirac operator on U .

Denote $i: U \hookrightarrow K$ the embedding. Let N be the normal bundle over U in K and $e \in H^2(U; \mathbb{Z})$ the Euler class of N . Clearly $i^*TK \cong TU \oplus N$.

Let E be a real vector bundle over K . Then i^*E is a real vector bundle over the spin manifold U . Let $\text{Ind}_2(i^*E)$ be the mod 2 index in the sense of AtiyahSinger [5] associated to i^*E , which is a spin cobordism invariant. Let $E_{\mathbb{C}}$ be the complexification of E .

We have the following analytic Rokhlin congruence formula.

Theorem 2.1 (Zhang [33, 34, 36]).

$$(2.1) \quad \left\langle \widehat{A}(TK) \exp\left(\frac{c}{2}\right) \text{ch}(E_{\mathbb{C}}), [K] \right\rangle \equiv \text{Ind}_2(D_U^{i^*E}) \pmod{2}$$

and

$$(2.2) \quad \left\langle \widehat{A}(TK) \text{ch}(E_{\mathbb{C}}), [K] \right\rangle \equiv \text{Ind}_2(D_U^{i^*E}) - \frac{1}{2} \left\langle \widehat{A}(TU) \text{ch}(i^*E_{\mathbb{C}}) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2}.$$

Combining (2.2) with Theorems 1.2 and 1.3, we have

Theorem 2.2. *If Z is a 12 dimensional closed smooth spin^c manifold and U is a characteristic submanifold, then the following identities hold,*

$$(2.3) \quad \left\langle \frac{C(x)[p - C(x)^2]}{12}, [Z] \right\rangle \\ = \left\langle 2\widehat{A}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + \widehat{A}(TZ) \text{ch}(T_{\mathbb{C}}Z) - 4\widehat{A}(TZ), [Z] \right\rangle \\ \equiv \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^N) - \frac{1}{2} \left\langle \widehat{A}(TU) \text{ch}(2i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} - 4) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2};$$

and

$$(2.4) \quad \left\langle \frac{\widetilde{C}(x)[\widetilde{p} + 6\lambda\widetilde{C}(x) - 4\widetilde{C}(x)^2]}{12}, [Z] \right\rangle \\ = \left\langle \widehat{A}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + \widehat{A}(TZ) \text{ch}(T_{\mathbb{C}}Z) + 244\widehat{A}(TZ), [Z] \right\rangle \\ \equiv \text{Ind}_2(D_U^{i^*V(x)}) + \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^N) \\ - \frac{1}{2} \left\langle \widehat{A}(TU) \text{ch}(i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} + 244) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2}.$$

2.1.2. Spin^{w_2} case. We call an $8k + 4$ dimensional closed smooth oriented manifold K a spin^{w_2} manifold if there exists a rank 2 nonorientable real vector bundle ξ such that $w_2(TK) = w_2(\xi)$. Such manifolds and the corresponding Rokhlin congruence have been studied in [34] and [37]. Let U be a *characteristic submanifold* of the spin^{w_2} structure, i.e., a nonorientable $8k + 2$ dimensional submanifold of K such that $[U] \in H_{8k+2}(K; \mathbb{Z}_2)$ is dual to $w_2(TK) \in H^2(K; \mathbb{Z}_2)$. U carries a canonically induced pin^- structure up to pin^- cobordism.

Denote $i : U \hookrightarrow K$ the embedding. Let N be the normal bundle over U in K and $e \in H^2(U, o(TU))$ the Euler class of N .

Let E be a real vector bundle over K . Then i^*E is a real vector bundle over the pin^- manifold U . Let $\text{Ind}_2^a(i^*E)$ be the mod 2 analytic index of the real vector bundle i^*E over U , which is defined via η -invariants (c.f. [37]). We have the following Rokhlin congruence formula.

Theorem 2.3 (Zhang (Theorem A.2 in [37])).

$$(2.5) \quad \left\langle \widehat{A}(TK) \text{ch}(E_{\mathbb{C}}), [K] \right\rangle \equiv \text{Ind}_2^a(i^*E) - \frac{1}{2} \left\langle \widehat{A}(TU) \text{ch}(i^*E_{\mathbb{C}}) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2}.$$

Combining (2.5) with Theorem 1.2 and Theorem 1.3, we have

Theorem 2.4. *If Z is a 12 dimensional closed smooth spin^{w_2} manifold and U is a characteristic submanifold, then the following identities hold,*

$$\begin{aligned}
(2.6) \quad & \left\langle \frac{C(x)[p - C(x)^2]}{12}, [Z] \right\rangle \\
&= \left\langle 2\hat{A}(TZ)\text{ch}(V_{\mathbb{C}}(x)) + \hat{A}(TZ)\text{ch}(T_{\mathbb{C}}Z) - 4\hat{A}(TZ), [Z] \right\rangle \\
&\equiv \text{Ind}_2^g(TU) + \text{Ind}_2^g(N) - \frac{1}{2} \left\langle \hat{A}(TU)\text{ch}(2i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} - 4) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2};
\end{aligned}$$

and

$$\begin{aligned}
(2.7) \quad & \left\langle \frac{\tilde{C}(x)[\tilde{p} + 6\lambda\tilde{C}(x) - 4\tilde{C}(x)^2]}{12}, [Z] \right\rangle \\
&= \left\langle \hat{A}(TZ)\text{ch}(V_{\mathbb{C}}(x)) + \hat{A}(TZ)\text{ch}(T_{\mathbb{C}}Z) + 244\hat{A}(TZ), [Z] \right\rangle \\
&\equiv \text{Ind}_2^g(i^*V(x)) + \text{Ind}_2^g(TU) + \text{Ind}_2^g(N) \\
&\quad - \frac{1}{2} \left\langle \hat{A}(TU)\text{ch}(i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} + 244) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2}.
\end{aligned}$$

2.2. New cubic forms on spin^c manifolds. Let Z be a 12 dimensional closed smooth spin^c manifold. Let ξ be the complex line bundle of the spin^c structure. We use $\xi_{\mathbb{R}}$ for the notation of ξ when it is viewed as an oriented real plane bundle. Let $c = c_1(\xi) \in H^2(Z; \mathbb{Z})$. Denote $\xi_{\mathbb{C}} = \xi_{\mathbb{R}} \otimes_{\mathbb{R}} \mathbb{C}$. Clearly $\xi_{\mathbb{C}} = \xi \oplus \bar{\xi}$.

Since Z is spin^c , $TZ \oplus \xi_{\mathbb{R}}$ is spin . By a result of McLaughlin (Lemma 2.2, [28]), there is a canonical class $\rho_c \in H^4(Z; \mathbb{Z})$ associated to the spin^c structure such that

$$2\rho_c = p_1(TZ \oplus \xi_{\mathbb{R}}) \in H^4(Z; \mathbb{Z}).$$

However,

$$\begin{aligned}
(2.8) \quad & p_1(TZ \oplus \xi_{\mathbb{R}}) \\
&= -c_2((TZ \oplus \xi_{\mathbb{R}}) \otimes_{\mathbb{R}} \mathbb{C}) \\
&= -c_2(T_{\mathbb{C}}Z) - c_2(\xi \oplus \bar{\xi}) - c_1(T_{\mathbb{C}}Z)c_1(\xi \oplus \bar{\xi}) \\
&= p_1(TZ) + c^2.
\end{aligned}$$

So

$$2\rho_c = p_1(TZ) + c^2.$$

Let

$$\lambda_c := \rho_c - 2c^2 \in H^4(Z; \mathbb{Z})$$

and

$$C_c(x) = \lambda_c + 2x \in H^4(Z; \mathbb{Z}).$$

Theorem 2.5. *There is a degree 8 integral class p_c such that*

$$\begin{aligned}
(2.9) \quad & p_c \equiv w_8 \pmod{2}, \\
& 8p_c = 4p_2 - p_1^2 - 6p_1c^2 + 39c^4.
\end{aligned}$$

Moreover, the following identity holds,

$$(2.10) \quad \left\langle \frac{C_c(x)[p_c - C_c(x)^2]}{24}, [Z] \right\rangle \\ = \left\langle \hat{A}(TZ)e^{c/2}\text{ch}(V_{\mathbb{C}}(x)) + \frac{1}{2}\hat{A}(TZ)e^{c/2}\text{ch}(T_{\mathbb{C}}Z) - \frac{1}{2}\hat{A}(TZ)e^{c/2}\text{ch}[\xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} - \xi_{\mathbb{C}} + 2], [Z] \right\rangle.$$

The existence of $p_c \in H^8(Z; \mathbb{Z})$ is proved in Theorem 5.1. Let

$$\tilde{p}_c = p_c - 3\lambda_c^2.$$

In Theorem 5.1, it is also shown that

$$(2.11) \quad \tilde{p}_c \equiv w_8 + w_4^2 + w_2^4 \pmod{2}, \\ 8\tilde{p}_c = 4p_2 - 7p_1^2 + 30p_1c^2 - 15c^4.$$

Let

$$\tilde{C}_c(x) = \lambda_c + x \in H^4(Z; \mathbb{Z}).$$

Theorem 2.6. *The following identity holds,*

$$(2.12) \quad \left\langle \frac{\tilde{C}_c(x)[\tilde{p}_c + 6\lambda_c\tilde{C}_c(x) - 4\tilde{C}_c(x)^2]}{12}, [Z] \right\rangle \\ = \left\langle \hat{A}(TZ)e^{c/2}\text{ch}(V_{\mathbb{C}}(x)) + \hat{A}(TZ)e^{c/2}\text{ch}(T_{\mathbb{C}}Z) + \hat{A}(TZ)e^{c/2}\text{ch}[-\xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} + \xi_{\mathbb{C}} + 246], [Z] \right\rangle.$$

These two theorems are consequences of the factorization formulas for degree 12 characteristic forms: (4.33) and (4.42) (proved in Section 4.2) and a direct computation of the degree 8 components in (4.33) and (4.42).

Let U be a characteristic submanifold of Z . Let N be the normal bundle over U in Z . Applying the Rokhlin congruence formula (2.1), we have

Theorem 2.7. *The following identities hold,*

$$(2.13) \quad \left\langle \frac{C_c(x)[p_c - C_c(x)^2]}{12}, [Z] \right\rangle \equiv \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^{N \otimes N}) \pmod{2};$$

and

$$(2.14) \quad \left\langle \frac{\tilde{C}_c(x)[\tilde{p}_c + 6\lambda_c\tilde{C}_c(x) - 4\tilde{C}_c(x)^2]}{12}, [Z] \right\rangle \equiv \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^{N \otimes N}) + \text{Ind}_2(D_U^{i^*V(x)}) \pmod{2}.$$

Proof. Clearly $i^*TZ \cong TU \oplus N$ and $i^*\xi_{\mathbb{R}} \cong N$.

Combining Theorem 2.5 with (2.1), we have, when mod 2,

$$(2.15) \quad \left\langle \frac{C_c(x)[p_c - C_c(x)^2]}{12}, [Z] \right\rangle \\ = \left\langle 2\hat{A}(TZ)e^{c/2}\text{ch}(V_{\mathbb{C}}(x)) + \hat{A}(TZ)e^{c/2}\text{ch}(T_{\mathbb{C}}Z) - \hat{A}(TZ)e^{c/2}[\xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} - \xi_{\mathbb{C}} + 2], [Z] \right\rangle \\ = 2\text{Ind}_2(D_U^{i^*V(x)}) + \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^N) + \text{Ind}_2(D_U^{N \otimes N}) + \text{Ind}_2(D_U^N) + 2\text{Ind}_2(D_U) \\ = \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^{N \otimes N}).$$

Combining Theorem 2.6 with (2.1), we have, when mod 2,

$$\begin{aligned}
(2.16) \quad & \left\langle \frac{\tilde{C}_c(x)[\tilde{p}_c + 6\lambda_c\tilde{C}_c(x) - 4\tilde{C}_c(x)^2]}{12}, [Z] \right\rangle \\
& = \left\langle \hat{A}(TZ)e^{c/2}\text{ch}(V_{\mathbb{C}}(x)) + \hat{A}(TZ)e^{c/2}\text{ch}(T_{\mathbb{C}}Z) + \hat{A}(TZ)e^{c/2}[-\xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} + \xi_{\mathbb{C}} + 246], [Z] \right\rangle \\
& = \text{Ind}_2(D_U^{i^*V(x)}) + \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^N) + \text{Ind}_2(D_U^{N \otimes N}) + \text{Ind}_2(D_U^N) + 246\text{Ind}_2(D_U) \\
& = \text{Ind}_2(D_U^{TU}) + \text{Ind}_2(D_U^{N \otimes N}) + \text{Ind}_2(D_U^{i^*V(x)}).
\end{aligned}$$

The desired formulas follow. \square

By subtracting the corresponding sides of (2.3) from (2.13), and (2.4) from (2.14) respectively, and then applying the Poincaré duality, we can show the following through direct computations.

Corollary 2.1. *Let e be the Euler class of N over U . The following identities hold,*

$$\begin{aligned}
(2.17) \quad & \frac{1}{64} \langle e \cdot \{24i^*C(x)^2 - (4p_1(TU) + 10e^2)i^*C(x) + p_1(TU)^2 - 4p_2(TU) + 6p_1(TU)e^2 - 21e^4\}, [U] \rangle \\
& \equiv \text{Ind}_2(D_U^N) + \text{Ind}_2(D_U^{N \otimes N}) + \frac{1}{2} \left\langle \hat{A}(TU)\text{ch}(2i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} - 4) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2};
\end{aligned}$$

and

$$\begin{aligned}
(2.18) \quad & \frac{1}{64} \langle e \cdot \{48i^*\tilde{C}(x)^2 - (28p_1(TU) + 10e^2)i^*\tilde{C}(x) + 7p_1(TU)^2 - 4p_2(TU) + 6p_1(TU)e^2 - 21e^4\}, [U] \rangle \\
& \equiv \text{Ind}_2(D_U^N) + \text{Ind}_2(D_U^{N \otimes N}) + \frac{1}{2} \left\langle \hat{A}(TU)\text{ch}(i^*V_{\mathbb{C}}(x) + T_{\mathbb{C}}U + N_{\mathbb{C}} + 244) \tanh\left(\frac{e}{4}\right), [U] \right\rangle \pmod{2}.
\end{aligned}$$

Remark 2.1. *Note that on the spin manifold U ,*

$$i^*C(x) = \frac{1}{2}p_1(TU) + 2i^*x, \quad i^*\tilde{C}(x) = \frac{1}{2}p_1(TU) + i^*x.$$

The left hand sides of (2.17) and (2.18) give very interesting quadratic forms on $H^4(U; \mathbb{Z})$ after being multiplied by 64.

Now suppose Z has boundary and let Y be the boundary of Z with the induced spin^c structure. Assume all the involved metrics and connections are of product structures near $\partial Z = Y$. Let D_Y^c be the Atiyah-Singer spin^c Dirac operator on Y .

Let

$$C_c(\tilde{x}) = \lambda_c(\nabla^{TZ}) + 2\tilde{x}$$

and

$$\tilde{C}_c(\tilde{x}) = \lambda_c(\nabla^{TZ}) + \tilde{x}.$$

As (4.33) and (4.42) hold on the level of forms, by the Atiyah-Patodi-Singer index theorem [2], we have the following formulas,

Theorem 2.8. *The following identities hold,*

$$(2.19) \quad \frac{1}{12} \int_Z C_c(\tilde{x})[p_c(\nabla^{TZ}) - C_c(\tilde{x})^2] \equiv 2\bar{\eta}(D_Y^{c, V_{\mathbb{C}}(x)}) + \bar{\eta}(D_Y^{c, T_{\mathbb{C}}Z}) - \bar{\eta}(D_Y^{c, \xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} - \xi_{\mathbb{C}} + 2}) \pmod{\mathbb{Z}};$$

and

$$\begin{aligned}
(2.20) \quad & \frac{1}{12} \int_Z \tilde{C}_c(\tilde{x})[\tilde{p}_c(\nabla^{TZ}) + 6\lambda_c(\nabla^{TZ})\tilde{C}_c(\tilde{x}) - 4\tilde{C}_c(\tilde{x})^2] \\
& \equiv \bar{\eta}(D_Y^{c, V_{\mathbb{C}}(x)}) + \bar{\eta}(D_Y^{c, T_{\mathbb{C}}Z}) + \bar{\eta}(D_Y^{c, -\xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} + \xi_{\mathbb{C}} + 246}) \pmod{\mathbb{Z}}.
\end{aligned}$$

3. CUBIC FORMS ON ORIENTABLE 12-MANIFOLDS

In this section, we give the Witten-Freed-Hopkins type anomaly cancellation formulas on 12 dimensional orientable manifolds, without assuming that the manifold is spin, spin^c or spin^{w2}.

Let Z be a 12 dimensional oriented smooth closed manifold. Let

$$(3.1) \quad \widehat{L}(x) = \frac{x}{\tanh \frac{x}{2}}$$

be the \widehat{L} -polynomial and $\widehat{L}(TZ)$ be the \widehat{L} -class of TZ (c.f. [26]). Let d_s be the signature operator on Z and W be a complex vector bundle over Z . Then by the Atiyah-Singer index theorem ([4])

$$\text{Ind}(d_s \otimes W) = \langle \widehat{L}(TZ) \text{ch}(W), [Z] \rangle.$$

For $x \in H^4(Z; \mathbb{Z})$, let

$$D(x) = -p_1 + 2x.$$

Theorem 3.1. *The following identity holds,*

$$(3.2) \quad \left\langle \frac{D(x)[4p_1^2 - 7p_2 - D(x)^2]}{6}, [Z] \right\rangle \\ = \frac{1}{32} \left\langle 2\widehat{L}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + 2\widehat{L}(TZ) \text{ch}(T_{\mathbb{C}}Z) + \widehat{L}(TZ) \text{ch}(\wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z)) - 4\widehat{L}(TZ), [Z] \right\rangle.$$

Let

$$\widetilde{D}(x) = -p_1 + x.$$

Theorem 3.2. *The following identity holds,*

$$(3.3) \quad \left\langle \frac{\widetilde{D}(x)[p_1^2 - 7p_2 - 6p_1\widetilde{D}(x) - 4\widetilde{D}(x)^2]}{3}, [Z] \right\rangle \\ = \frac{1}{16} \left\langle \widehat{L}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + 2\widehat{L}(TZ) \text{ch}(T_{\mathbb{C}}Z) + \widehat{L}(TZ) \text{ch}(\wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z)) + 244\widehat{L}(TZ), [Z] \right\rangle.$$

These two theorems are consequences of the factorization formulas for degree 12 characteristic forms: (4.59) and (4.69) (proved in Subsection 4.3) and direct computations of the degree 8 components in (4.59) and (4.69).

Now suppose Z has boundary and let Y be the boundary of Z with the induced orientation. Assume all the involved metrics and connections are of product structures near $\partial Z = Y$. Let B_Y be the signature operator on Y .

Denote

$$D(\check{x}) = -p_1(\nabla^{TZ}) + 2\check{x}$$

and

$$\widetilde{D}(\check{x}) = -p_1(\nabla^{TZ}) + \check{x}.$$

As (4.59) and (4.69) hold on the level of forms, by the Atiyah-Patodi-Singer index theorem [2], we have

Theorem 3.3. *The following identities hold,*

$$(3.4) \quad \begin{aligned} & \frac{16}{3} \int_Z D(\tilde{x}) [4p_1(\nabla^{TZ})^2 - 7p_2(\nabla^{TZ}) - D(\tilde{x})^2] \\ & \equiv 2\bar{\eta}(B_Y^{V_C(x)}) + 2\bar{\eta}(B_Y^{T_C Z}) + \bar{\eta}(B_Y^{\wedge^2(T_C Z) - S^2(T_C Z)}) - 4\bar{\eta}(B_Y) \pmod{\mathbb{Z}}; \end{aligned}$$

and

$$(3.5) \quad \begin{aligned} & \frac{16}{3} \int_Z \tilde{D}(\tilde{x}) [p_1(\nabla^{TZ})^2 - 7p_2(\nabla^{TZ}) - 6p_1(\nabla^{TZ})\tilde{D}(\tilde{x}) - 4\tilde{D}(\tilde{x})^2] \\ & \equiv \bar{\eta}(B_Y^{V_C(x)}) + 2\bar{\eta}(B_Y^{T_C Z}) + \bar{\eta}(B_Y^{\wedge^2(T_C Z) - S^2(T_C Z)}) + 244\bar{\eta}(B_Y) \pmod{\mathbb{Z}}. \end{aligned}$$

4. PROOFS

The purpose of this section is to give proofs to Theorem 2.5, Theorem 2.6, Theorem 3.1 and Theorem 3.2. In order to conduct the proofs, we first briefly review some basic materials on the representation theory of affine E_8 following [24] (see also [25]).

4.1. The basic representation of affine E_8 . Let \mathfrak{g} be the Lie algebra of E_8 . Let $\langle \cdot, \cdot \rangle$ be the Killing form on \mathfrak{g} . Let $\tilde{\mathfrak{g}}$ be the affine Lie algebra corresponding to \mathfrak{g} defined by

$$\tilde{\mathfrak{g}} = \mathbb{C}[t, t^{-1}] \otimes \mathfrak{g} \oplus \mathbb{C}c,$$

with the bracket

$$[P(t) \otimes x + \lambda c, Q(t) \otimes y + \mu c] = P(t)Q(t) \otimes [x, y] + \langle x, y \rangle \text{Res}_{t=0} \left(\frac{dP(t)}{dt} Q(t) \right) c.$$

Let $\hat{\mathfrak{g}}$ be the affine Kac-Moody algebra obtained from $\tilde{\mathfrak{g}}$ by adding a derivation $t \frac{d}{dt}$ which operates on $\mathbb{C}[t, t^{-1}] \otimes \mathfrak{g}$ in an obvious way and sends c to 0.

The basic representation $V(\Lambda_0)$ is the $\hat{\mathfrak{g}}$ -module defined by the property that there is a nonzero vector v_0 (highest weight vector) in $V(\Lambda_0)$ such that $cv_0 = v_0, (\mathbb{C}[t] \oplus \mathbb{C}t \frac{d}{dt})v_0 = 0$. Setting $V_k := \{v \in V(\Lambda_0) | t \frac{d}{dt} v = -kv\}$ gives a \mathbb{Z}_+ -grading by finite spaces. Since $[g, d] = 0$, each V_k is a representation ρ_k of \mathfrak{g} . Moreover, ρ_1 is the adjoint representation of E_8 .

Let $q = e^{2\pi\sqrt{-1}\tau}, \tau \in \mathbb{H}$. Fix a basis for the Cartan subalgebra and let $\{z_i\}_{i=1}^8$ be the corresponding coordinates. The character of the basic representation is given by

$$\text{ch}(z_1, z_2, \dots, z_8, \tau) := \sum_{k=0}^{\infty} (\text{ch} V_k)(z_1, z_2, \dots, z_8) q^k = \varphi(\tau)^{-8} \Theta_{\mathfrak{g}}(z_1, z_2, \dots, z_8, \tau),$$

where $\varphi(\tau) = \prod_{n=1}^{\infty} (1 - q^n)$ so that $\eta(\tau) = q^{1/24} \varphi(\tau)$ is the Dedekind η function; $\Theta_{\mathfrak{g}}(z_1, z_2, \dots, z_8, \tau)$ is the theta function defined on the root lattice Q by

$$\Theta_{\mathfrak{g}}(z_1, z_2, \dots, z_8, \tau) = \sum_{\gamma \in Q} q^{|\gamma|^2/2} e^{2\pi\sqrt{-1}\gamma(\vec{z})},$$

where $\vec{z} = (z_1, z_2, \dots, z_8)$.

It was proved in [12] that there is a basis for the E_8 root lattice such that

$$(4.1) \quad \Theta_{\mathfrak{g}}(z_1, \dots, z_8, \tau) = \frac{1}{2} \left(\prod_{l=1}^8 \theta(z_l, \tau) + \prod_{l=1}^8 \theta_1(z_l, \tau) + \prod_{l=1}^8 \theta_2(z_l, \tau) + \prod_{l=1}^8 \theta_3(z_l, \tau) \right),$$

where θ and θ_i ($i = 1, 2, 3$) are the Jacobi theta functions (c.f. [8] and [17]):

$$(4.2) \quad \theta(v, \tau) = 2q^{1/8} \sin(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^j)(1 - e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(4.3) \quad \theta_1(v, \tau) = 2q^{1/8} \cos(\pi v) \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^j)(1 + e^{-2\pi\sqrt{-1}v} q^j)],$$

$$(4.4) \quad \theta_2(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 - e^{2\pi\sqrt{-1}v} q^{j-1/2})(1 - e^{-2\pi\sqrt{-1}v} q^{j-1/2})],$$

$$(4.5) \quad \theta_3(v, \tau) = \prod_{j=1}^{\infty} [(1 - q^j)(1 + e^{2\pi\sqrt{-1}v} q^{j-1/2})(1 + e^{-2\pi\sqrt{-1}v} q^{j-1/2})].$$

The theta functions satisfy the the following transformation laws (c.f. [8]),

$$(4.6) \quad \theta(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta(v, \tau), \quad \theta(v, -1/\tau) = \frac{1}{\sqrt{-1}} \left(\frac{\tau}{\sqrt{-1}} \right)^{1/2} e^{\pi\sqrt{-1}\tau v^2} \theta(\tau v, \tau);$$

$$(4.7) \quad \theta_1(v, \tau + 1) = e^{\frac{\pi\sqrt{-1}}{4}} \theta_1(v, \tau), \quad \theta_1(v, -1/\tau) = \left(\frac{\tau}{\sqrt{-1}} \right)^{1/2} e^{\pi\sqrt{-1}\tau v^2} \theta_2(\tau v, \tau);$$

$$(4.8) \quad \theta_2(v, \tau + 1) = \theta_3(v, \tau), \quad \theta_2(v, -1/\tau) = \left(\frac{\tau}{\sqrt{-1}} \right)^{1/2} e^{\pi\sqrt{-1}\tau v^2} \theta_1(\tau v, \tau);$$

$$(4.9) \quad \theta_3(v, \tau + 1) = \theta_2(v, \tau), \quad \theta_3(v, -1/\tau) = \left(\frac{\tau}{\sqrt{-1}} \right)^{1/2} e^{\pi\sqrt{-1}\tau v^2} \theta_3(\tau v, \tau).$$

4.2. Proof of Theorem 2.5 and Theorem 2.6. Now we are ready to give the proofs.

4.2.1. *Proof of Theorem 2.5.* The proof of the statement (2.9) about p_c can be found in Theorem 5.1.

For the principal E_8 bundles P_i , $i = 1, 2$, consider the associated bundles

$$\mathcal{V}_i = \sum_{k=0}^{\infty} (P_i \times_{\rho_k} V_k) q^k \in K(Z)[[q]].$$

Let $W_i = P_i \times_{\rho_1} V_1$, $i = 1, 2$, be the complex vector bundles associated to the adjoint representation ρ_1 .

By the knowledge reviewed in Section 4.1, we see that there are formally two forms y_l^i ($i = 1, 2$ and $1 \leq l \leq 8$) on Z such that

$$(4.10) \quad \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) = \frac{1}{2} \left(\prod_{l=1}^8 \theta(y_l^i, \tau) + \prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right).$$

Since $\theta(z, \tau)$ is an odd function about z , one can see that up to degree 12, the term $\prod_{l=1}^8 \theta(y_l^i, \tau)$ can be dropped and therefore we have

$$(4.11) \quad \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) = \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right).$$

Since $\theta_1(z, \tau), \theta_2(z, \tau)$ and $\theta_3(z, \tau)$ are all even functions about z , the right hand side of the above equality only contains even powers of y_j^i 's. Therefore $\text{ch}(W_i)$ only consists of forms of degrees divisible by 4 (this is actually a basic fact about E_8). So

$$(4.12) \quad \text{ch}(\mathcal{V}_i) = 1 + \text{ch}(W_i)q + \cdots = 1 + (248 - c_2(W_i) + \cdots)q + \cdots.$$

On the other hand,

$$(4.13) \quad \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) = 1 + \left(240 + 30 \sum_{l=1}^8 (y_l^i)^2 + \dots \right) q + O(q^2).$$

From (4.11), (4.12) and (4.13), we have

$$(4.14) \quad \sum_{l=1}^8 (y_l^i)^2 = -\frac{1}{30} c_2(W_i).$$

Let TZ be the tangent bundle of Z and $T_{\mathbb{C}}Z$ be its complexification. Let ξ be a rank two real oriented Euclidean vector bundle over Z carrying a Euclidean connection ∇^ξ . Let $c = e(\xi, \nabla^\xi)$ be the Euler form canonically associated to ∇^ξ .

If E is a complex vector bundle over Z , set $\tilde{E} = E - \mathbb{C}^{\text{rk}(E)}$. Recall that for an indeterminate t ,

$$(4.15) \quad \Lambda_t(E) = \mathbb{C}|_Z + tE + t^2 \wedge^2(E) + \dots, \quad S_t(E) = \mathbb{C}|_Z + tE + t^2 S^2(E) + \dots,$$

are the total exterior and symmetric powers of E respectively. The following relations between these two operations hold (c.f. [3]),

$$(4.16) \quad S_t(E) = \frac{1}{\Lambda_{-t}(E)}, \quad \Lambda_t(E - F) = \frac{\Lambda_t(E)}{\Lambda_t(F)}.$$

Following [9], set

$$\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}}) := \left(\bigotimes_{m=1}^{\infty} S_{q^m}(\widetilde{T_{\mathbb{C}}Z}) \right) \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\tilde{\xi}_{\mathbb{C}}) \right) \otimes \left(\bigotimes_{u=1}^{\infty} \Lambda_{-q^{u-1/2}}(\tilde{\xi}_{\mathbb{C}}) \right) \otimes \left(\bigotimes_{v=1}^{\infty} \Lambda_{q^{v-1/2}}(\tilde{\xi}_{\mathbb{C}}) \right) \in K(Z)[[q]],$$

where $\xi_{\mathbb{C}}$ is the complexification of ξ .

Clearly, $\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})$ admits a formal Fourier expansion in q as

$$(4.17) \quad \Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}}) = \mathbb{C} + B_1 q + B_2 q^2 \dots,$$

where the B_j 's are elements in the semi-group formally generated by complex vector bundles over Z . Moreover, they carry canonically induced connections denoted by ∇^{B_j} 's. Let ∇^{Θ} be the induced connection with q -coefficients on Θ .

For $1 \leq i, j \leq 2$, set

$$(4.18) \quad \mathcal{Q}(P_i, P_j, \xi, \tau) := \left\{ e^{\frac{1}{24} E_2(\tau) (p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \hat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \right\}^{(12)}.$$

Here

$$(4.19) \quad E_2(\tau) = 1 - 24 \sum_{n=1}^{\infty} \left(\sum_{d|n} d \right) q^n$$

is the Eisenstein series. Unlike the other Eisenstein series $E_{2k}(\tau)$, $k > 1$, $E_2(\tau)$ is not a modular form over $SL(2, \mathbb{Z})$, instead $E_2(\tau)$ is a quasimodular form over $SL(2, \mathbb{Z})$, satisfying:

$$(4.20) \quad E_2\left(\frac{a\tau + b}{c\tau + d}\right) = (c\tau + d)^2 E_2(\tau) - \frac{6\sqrt{-1}c(c\tau + d)}{\pi}.$$

In particular, we have

$$(4.21) \quad E_2(\tau + 1) = E_2(\tau),$$

$$(4.22) \quad E_2\left(-\frac{1}{\tau}\right) = \tau^2 E_2(\tau) - \frac{6\sqrt{-1}\tau}{\pi}$$

(c.f. Chap 2.3 in [7]).

Lemma 4.1. $\mathcal{Q}(P_i, P_j, \xi, \tau)$ is a modular form of weight 14 over $SL(2; \mathbb{Z})$.

Proof. Let $\{\pm 2\pi\sqrt{-1}x_k\} (1 \leq k \leq 6)$ be the formal Chern roots for $(T_{\mathbb{C}}Z, \nabla^{T_{\mathbb{C}}Z})$. Let $c = e(\xi, \nabla^\xi) = 2\pi\sqrt{-1}u$. One has

$$(4.23) \quad \begin{aligned} & \mathcal{Q}(P_i, P_j, \xi, \tau) \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \right\}^{(12)} \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \left(\prod_{k=1}^6 \left(x_k \frac{\theta'(0, \tau)}{\theta(x_k, \tau)} \right) \right) \frac{\theta_1(u, \tau)}{\theta_1(0, \tau)} \frac{\theta_2(u, \tau)}{\theta_2(0, \tau)} \frac{\theta_3(u, \tau)}{\theta_3(0, \tau)} \right. \\ & \quad \left. \cdot \frac{1}{4} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \left(\prod_{l=1}^8 \theta_1(y_l^j, \tau) + \prod_{l=1}^8 \theta_2(y_l^j, \tau) + \prod_{l=1}^8 \theta_3(y_l^j, \tau) \right) \right\}^{(12)}. \end{aligned}$$

Then we can perform the transformation laws (4.6)-(4.9) for the theta functions and the transformation laws (4.21), (4.22) for $E_2(\tau)$ to show that $\mathcal{Q}(P_i, P_j, \xi, \tau)$ is a modular form of weight 14 over $SL(2; \mathbb{Z})$. \square

Expanding the q -series, using (4.12), (4.17) and (4.19), we have

$$(4.24) \quad \begin{aligned} & e^{\frac{1}{24}E_2(\tau)(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \\ &= \left(e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \right. \\ & \quad \left. - e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) q + O(q^2) \right) \\ & \quad \cdot \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\mathbb{C} + B_1q + O(q^2))(1 - 16q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2))(1 + \text{ch}(W_j)q + O(q^2)) \\ &= e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \\ & \quad + q \left(e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(B_1 - 16 + W_i + W_j) \right. \\ & \quad \left. - e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}(c_2(W_i)+c_2(W_j)))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right) \\ & \quad + O(q^2). \end{aligned}$$

It is well known (c.f. Chap 2.1 in [7]) that modular forms over $SL(2; \mathbb{Z})$ can be expressed as polynomials of the Eisenstein series $E_4(\tau)$, $E_6(\tau)$, where

$$(4.25) \quad E_4(\tau) = 1 + 240q + 2160q^2 + 6720q^3 + \dots,$$

$$(4.26) \quad E_6(\tau) = 1 - 504q - 16632q^2 - 122976q^3 + \dots.$$

Their weights are 4 and 6 respectively.

Since the weight of the modular form $\mathcal{Q}(P_i, P_j, \xi, \tau)$ is 14 and the space of modular forms of weight 14 over $SL(2, \mathbb{Z})$ is 1-dimensional and spanned by $E_4^2(\tau)E_6(\tau)$ (c.f. Chap 2.1 in [7]), it must be a multiple of

$$(4.27) \quad E_4(\tau)^2 E_6(\tau) = 1 - 24q + \dots$$

So from (4.24) and (4.27), we have

$$(4.28) \quad \begin{aligned} & \left\{ e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(B_1 - 16 + W_i + W_j) \right\}^{(12)} \\ & - \left\{ e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(12)} \\ = & - 24 \left\{ e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(12)}. \end{aligned}$$

Therefore

$$(4.29) \quad \begin{aligned} & \left\{ \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + W_j + B_1 + 8) \right\}^{(12)} \\ = & \left(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \\ & \cdot \left\{ - \frac{e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + W_j + B_1 + 8) \right. \\ & \left. + e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(8)}. \end{aligned}$$

To find B_1 , we have

$$(4.30) \quad \begin{aligned} & \Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}}) \\ = & \left(\bigotimes_{m=1}^{\infty} S_{qm}(\widetilde{T_{\mathbb{C}}Z}) \right) \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{qn}(\widetilde{\xi_{\mathbb{C}}}) \right) \otimes \left(\bigotimes_{u=1}^{\infty} \Lambda_{-qu-1/2}(\widetilde{\xi_{\mathbb{C}}}) \right) \otimes \left(\bigotimes_{v=1}^{\infty} \Lambda_{qv-1/2}(\widetilde{\xi_{\mathbb{C}}}) \right) \\ = & 1 + (T_{\mathbb{C}}Z - 12 - 3\widetilde{\xi_{\mathbb{C}}} - \widetilde{\xi_{\mathbb{C}}} \otimes \widetilde{\xi_{\mathbb{C}}})q + O(q^2). \end{aligned}$$

So

$$(4.31) \quad B_1 = T_{\mathbb{C}}Z - 12 - 3\widetilde{\xi_{\mathbb{C}}} - \widetilde{\xi_{\mathbb{C}}} \otimes \widetilde{\xi_{\mathbb{C}}}.$$

Plugging B_1 into (4.29), we have

$$(4.32) \quad \begin{aligned} & \left\{ \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + W_j + T_{\mathbb{C}}Z - 4 - 3\widetilde{\xi_{\mathbb{C}}} - \widetilde{\xi_{\mathbb{C}}} \otimes \widetilde{\xi_{\mathbb{C}}}) \right\}^{(12)} \\ = & \left(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \\ & \cdot \left\{ - \frac{e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + W_j + T_{\mathbb{C}}Z - 4 - 3\widetilde{\xi_{\mathbb{C}}} - \widetilde{\xi_{\mathbb{C}}} \otimes \widetilde{\xi_{\mathbb{C}}}) \right. \\ & \left. + e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(8)}. \end{aligned}$$

Since $\text{ch}(W_i), \text{ch}(W_j)$ only contribute degree $4l$ forms, we can replace $\cosh\left(\frac{c}{2}\right)$ by $e^{\frac{c}{2}}$. Then in (4.32), putting $W_1 = W_2 = V_{\mathbb{C}}(x)$, we get

$$(4.33) \quad \begin{aligned} & \left\{ \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(V_{\mathbb{C}}(x)) + \frac{1}{2} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(T_{\mathbb{C}}Z) - \frac{1}{2} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(4 + 3\widetilde{\xi}_{\mathbb{C}} + \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}) \right\}^{(12)} \\ &= \left(\frac{p_1(TZ) - 3c^2}{2} + \frac{1}{30} c_2(V_{\mathbb{C}}(x)) \right) \\ & \cdot \left\{ -\frac{e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x))} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(\mathfrak{A}) + e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} \widehat{A}(TZ) e^{\frac{c}{2}} \right\}^{(8)}, \end{aligned}$$

where

$$\mathfrak{A} = 2V_{\mathbb{C}}(x) + T_{\mathbb{C}}Z - 4 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}.$$

It is not hard to check that $4 + 3\widetilde{\xi}_{\mathbb{C}} + \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}} = \xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} - \xi_{\mathbb{C}} + 2$ and

$$\frac{p_1(TZ) - 3c^2}{2} + \frac{1}{30} c_2(V_{\mathbb{C}}(x)) = \lambda_c + 2x = C_c(x).$$

A direct computation shows that the 8-form in the right hand side of (4.33) verifies

$$(4.34) \quad \begin{aligned} & \left\{ -\frac{e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x))} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(\mathfrak{A}) + e^{\frac{1}{24}(p_1(TZ) - 3c^2 + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} \widehat{A}(TZ) e^{\frac{c}{2}} \right\}^{(8)} \\ &= \frac{p_c - C_c(x)^2}{24}. \end{aligned}$$

We therefore get (2.10), and have completed the proof of Theorem 2.5.

4.2.2. *Proof of Theorem 2.6.* For each i , set

$$(4.35) \quad \begin{aligned} & \mathcal{R}(P_i, \xi, \tau) \\ & := \left\{ e^{\frac{1}{24}E_2(\tau)(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)}. \end{aligned}$$

Lemma 4.2. $\mathcal{R}(P_i, \xi, \tau)$ is a modular form of weight 10 over $SL(2; \mathbb{Z})$.

Proof. This can be similarly proved as Lemma 4.1 by seeing that

$$(4.36) \quad \begin{aligned} & \mathcal{R}(P_i, \xi, \tau) \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)} \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i))} \left(\prod_{l=1}^6 \left(x_l \frac{\theta'(0, \tau)}{\theta(x_l, \tau)} \right) \right) \frac{\theta_1(u, \tau)}{\theta_1(0, \tau)} \frac{\theta_2(u, \tau)}{\theta_2(0, \tau)} \frac{\theta_3(u, \tau)}{\theta_3(0, \tau)} \right. \\ & \quad \left. \cdot \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(12)}, \end{aligned}$$

and then we can perform the transformation laws (4.6)-(4.9) for the theta functions and the transformation laws (4.21), (4.22) for $E_2(\tau)$ to show that $\mathcal{R}(P_i, \xi, \tau)$ is a modular form of weight 10 over $SL(2; \mathbb{Z})$. \square

Similar to the proof of Theorem 2.5, expanding the q -series, using (4.12), (4.17) and (4.19), we have

$$\begin{aligned}
& e^{\frac{1}{24}E_2(\tau)(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\Theta(T_{\mathbb{C}}Z, \xi_{\mathbb{C}})) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \\
&= \left(e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \right. \\
&\quad \left. - e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i) \right) q + O(q^2) \right) \\
&\quad \cdot \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(\mathbb{C} + B_1q + O(q^2))(1 - 8q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2)) \\
(4.37) \quad &= e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \\
&\quad + q \left(e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(B_1 - 8 + W_i) \right. \\
&\quad \left. - e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i) \right) \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right) \\
&\quad + O(q^2).
\end{aligned}$$

However modular form of weight 10 must be a multiple of (c.f. Chap 2.1 in [7])

$$(4.38) \quad E_4(\tau)E_6(\tau) = 1 - 264q + \dots$$

So from (4.37) and (4.38), we have

$$\begin{aligned}
& \left\{ e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(B_1 - 8 + W_i) \right\}^{(12)} \\
(4.39) \quad & - \left\{ e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i) \right) \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(12)} \\
&= -264 \left\{ e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(12)}.
\end{aligned}$$

Therefore

$$\begin{aligned}
& \left\{ \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + B_1 + 256) \right\}^{(12)} \\
(4.40) \quad &= \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i) \right) \\
&\quad \cdot \left\{ - \frac{e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i)} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + B_1 + 256) \right. \\
&\quad \left. + e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(8)}.
\end{aligned}$$

Plugging in B_1 (see (4.31)), we have

$$\begin{aligned}
& \left\{ \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + T_{\mathbb{C}}Z + 244 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}) \right\}^{(12)} \\
&= \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i) \right) \\
(4.41) \quad & \cdot \left\{ -\frac{e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}c_2(W_i)} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \text{ch}(W_i + T_{\mathbb{C}}Z + 244 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}) \right. \\
& \quad \left. + e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(W_i))} \widehat{A}(TZ) \cosh\left(\frac{c}{2}\right) \right\}^{(8)}.
\end{aligned}$$

Since $\text{ch}(W_i)$ only contributes degree $4l$ forms, we can replace $\cosh\left(\frac{c}{2}\right)$ by $e^{\frac{c}{2}}$. Taking $W_i = V_{\mathbb{C}}(x)$, we have

$$\begin{aligned}
(4.42) \quad & \left\{ \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(V_{\mathbb{C}}(x)) + \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(T_{\mathbb{C}}Z) + \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(244 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}) \right\}^{(12)} \\
&= \left(p_1(TZ) - 3c^2 + \frac{1}{30}c_2(V_{\mathbb{C}}(x)) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(V_{\mathbb{C}}(x)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}c_2(V_{\mathbb{C}}(x))} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(\mathfrak{B}) + e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(V_{\mathbb{C}}(x)))} \widehat{A}(TZ) e^{\frac{c}{2}} \right\}^{(8)},
\end{aligned}$$

where

$$\mathfrak{B} = V_{\mathbb{C}}(x) + T_{\mathbb{C}}Z + 244 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}}.$$

It is not hard to check that $244 - 3\widetilde{\xi}_{\mathbb{C}} - \widetilde{\xi}_{\mathbb{C}} \otimes \widetilde{\xi}_{\mathbb{C}} = 246 - \xi_{\mathbb{C}} \otimes \xi_{\mathbb{C}} + \xi_{\mathbb{C}}$ and

$$p_1(TZ) - 3c^2 + \frac{1}{30}c_2(V_{\mathbb{C}}(x)) = 2\lambda_c + 2x = 2\widetilde{C}_c(x).$$

A direct computation shows that the 8-form in the right hand side of (4.42) verifies

$$\begin{aligned}
& \left\{ -\frac{e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(V_{\mathbb{C}}(x)))} - 1}{p_1(TZ) - 3c^2 + \frac{1}{30}c_2(V_{\mathbb{C}}(x))} \widehat{A}(TZ) e^{\frac{c}{2}} \text{ch}(\mathfrak{E}) + e^{\frac{1}{24}(p_1(TZ)-3c^2+\frac{1}{30}c_2(V_{\mathbb{C}}(x)))} \widehat{A}(TZ) e^{\frac{c}{2}} \right\}^{(8)} \\
&= \frac{\widetilde{p}_c + 6\lambda_c \widetilde{C}_c(x) - 4\widetilde{C}_c(x)^2}{24}.
\end{aligned}$$

We therefore get (2.12), and have completed the proof of Theorem 2.6.

4.3. Proof of Theorem 3.1 and Theorem 3.2.

4.3.1. *Proof of Theorem 3.1.* Following [9], set

$$(4.43) \quad \Theta_1(T_{\mathbb{C}}Z) := \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\widetilde{T_{\mathbb{C}}Z}) \right) \in K(Z)[[q]],$$

$$(4.44) \quad \Theta_2(T_{\mathbb{C}}Z) := \left(\bigotimes_{n=1}^{\infty} \Lambda_{-q^{n-1/2}}(\widetilde{T_{\mathbb{C}}Z}) \right) \in K(Z)[[q^{1/2}]],$$

$$(4.45) \quad \Theta_3(T_{\mathbb{C}}Z) := \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^{n-1/2}}(\widetilde{T_{\mathbb{C}}Z}) \right) \in K(Z)[[q^{1/2}]].$$

Construct ([27])

$$(4.46) \quad \Phi(T_{\mathbb{C}}Z) = \Theta(T_{\mathbb{C}}Z) \otimes \Theta_1(T_{\mathbb{C}}Z) \otimes \Theta_2(T_{\mathbb{C}}Z) \otimes \Theta_3(T_{\mathbb{C}}Z) \in K(Z)[[q]]$$

and define

$$(4.47) \quad \mathcal{W}_{\widehat{L}}(TZ) = e^{-\frac{1}{12}E_2(\tau) \cdot p_1(TZ)} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \in H^{4*}(TZ, \mathbb{Q}).$$

We call $\mathcal{W}_{\widehat{L}}(TZ)$ the \widehat{L} -Witten class of TZ .

Perform the formal Fourier expansion in q as

$$(4.48) \quad \Phi(T_{\mathbb{C}}Z) = \Theta(T_{\mathbb{C}}Z) \otimes \Theta_1(T_{\mathbb{C}}Z) \otimes \Theta_2(T_{\mathbb{C}}Z) \otimes \Theta_3(T_{\mathbb{C}}Z) = \mathbb{C} + D_1q + D_2q^2 \cdots,$$

where the D_j 's are elements in the semi-group formally generated by complex vector bundles over Z . Moreover, they carry canonically induced connections denoted by ∇^{D_j} 's. Let ∇^{Φ} be the induced connection with q -coefficients on Φ .

For $1 \leq i, j \leq 2$, construct the twisted \widehat{L} -Witten class

$$(4.49) \quad e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \in H^{4*}(Z, \mathbb{Q})$$

and denote

$$(4.50) \quad \begin{aligned} & \mathcal{Q}_L(P_i, P_j, \tau) \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \right\}^{(12)}. \end{aligned}$$

Lemma 4.3. $\mathcal{Q}_L(P_i, P_j, \tau)$ is a modular form of weight 14 over $SL(2; \mathbb{Z})$.

Proof. Let $\{\pm 2\pi\sqrt{-1}x_k\} (1 \leq k \leq 6)$ be the formal Chern roots for $(T_{\mathbb{C}}Z, \nabla^{T_{\mathbb{C}}Z})$. One has

$$(4.51) \quad \begin{aligned} & \mathcal{Q}_L(P_i, P_j, \tau) \\ &= \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \right\}^{(12)} \\ &= 2^6 \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \left(\prod_{k=1}^6 x_k \frac{\theta'(0, \tau)}{\theta(x_k, \tau)} \frac{\theta_1(x_k, \tau)}{\theta_1(0, \tau)} \frac{\theta_2(x_k, \tau)}{\theta_2(0, \tau)} \frac{\theta_3(x_k, \tau)}{\theta_3(0, \tau)} \right) \right. \\ & \quad \left. \cdot \frac{1}{4} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \left(\prod_{l=1}^8 \theta_1(y_l^j, \tau) + \prod_{l=1}^8 \theta_2(y_l^j, \tau) + \prod_{l=1}^8 \theta_3(y_l^j, \tau) \right) \right\}^{(12)}. \end{aligned}$$

Then we can perform the transformation laws (4.6)-(4.9) for the theta functions and the transformation laws (4.21), (4.22) for $E_2(\tau)$ to show that $\mathcal{Q}_L(P_i, P_j, \tau)$ is a modular form of weight 14 over $SL(2; \mathbb{Z})$. \square

Expanding the q -series, using (4.12), (4.19) and (4.48), we have

$$\begin{aligned}
(4.52) \quad & e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^{16} \text{ch}(\mathcal{V}_i) \text{ch}(\mathcal{V}_j) \\
& = \left(e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \right. \\
& \quad \left. - e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \left(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) q + O(q^2) \right) \\
& \quad \cdot \widehat{L}(TZ) \text{ch}(\mathbb{C} + D_1 q + O(q^2))(1 - 16q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2))(1 + \text{ch}(W_j)q + O(q^2)) \\
& = e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \\
& \quad + q \left(e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(D_1 - 16 + W_i + W_j) \right. \\
& \quad \left. - e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \left(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \widehat{L}(TZ) \right) + O(q^2).
\end{aligned}$$

Since the weight of the modular form $\mathcal{Q}(P_i, P_j, \tau)$ is 14, it must be a multiple of

$$(4.53) \quad E_4(\tau)^2 E_6(\tau) = 1 - 24q + \dots$$

So from (4.52) and (4.53), we have

$$\begin{aligned}
(4.54) \quad & \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \text{ch}(D_1 - 16 + W_i + W_j) \right\}^{(12)} \\
& \quad - \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \left(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \widehat{L}(TZ) \right\}^{(12)} \\
& = -24 \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \right\}^{(12)}.
\end{aligned}$$

Therefore

$$\begin{aligned}
(4.55) \quad & \left\{ \widehat{L}(TZ) \text{ch}(W_i + W_j + D_1 + 8) \right\}^{(12)} \\
& = \left(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \\
& \quad \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} - 1}{-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j))} \widehat{L}(TZ) \text{ch}(W_i + W_j + D_1 + 8) \right. \\
& \quad \left. + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \right\}^{(8)}.
\end{aligned}$$

To find D_1 , we have

$$\begin{aligned}
(4.56) \quad & \Phi(T_{\mathbb{C}}Z) \\
& = \left(\bigotimes_{m=1}^{\infty} S_{q^m}(\widetilde{T_{\mathbb{C}}Z}) \right) \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^n}(\widetilde{T_{\mathbb{C}}Z}) \right) \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{-q^{n-1/2}}(\widetilde{T_{\mathbb{C}}Z}) \right) \otimes \left(\bigotimes_{n=1}^{\infty} \Lambda_{q^{n-1/2}}(\widetilde{T_{\mathbb{C}}Z}) \right) \\
& = 1 + (2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) - 12)q + O(q^2).
\end{aligned}$$

So

$$(4.57) \quad D_1 = 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) - 12.$$

Plugging D_1 into (4.55), we have

$$\begin{aligned}
(4.58) \quad & \left\{ \widehat{L}(TZ) \text{ch}(W_i + W_j + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) - 4) \right\}^{(12)} \\
&= \left(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} - 1}{-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j))} \widehat{L}(TZ) \text{ch}(W_i + W_j + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) - 4) + \right. \\
& \quad \left. + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}(c_2(W_i) + c_2(W_j)))} \widehat{L}(TZ) \right\}^{(8)}.
\end{aligned}$$

Putting $W_1 = W_2 = V_{\mathbb{C}}(x)$, we get

$$\begin{aligned}
(4.59) \quad & \left\langle 2\widehat{L}(TZ) \text{ch}(V_{\mathbb{C}}(x)) + 2\widehat{L}(TZ) \text{ch}(T_{\mathbb{C}}Z) + \widehat{L}(TZ) \text{ch}(\wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z)) - 4\widehat{L}(TZ), [Z] \right\rangle \\
&= \left(-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} - 1}{-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x))} \widehat{L}(TZ) \text{ch}(\mathfrak{C}) + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} \widehat{L}(TZ) \right\}^{(8)},
\end{aligned}$$

where

$$\mathfrak{C} = 2V_{\mathbb{C}}(x) + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) - 4.$$

Note that the 4-form

$$-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)) = 2D(x).$$

A direct computation shows that the 8-form in the right hand side of (4.59) verifies

$$\begin{aligned}
(4.60) \quad & \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} - 1}{-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x))} \widehat{L}(TZ) \text{ch}(\mathfrak{C}) + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{15}c_2(V_{\mathbb{C}}(x)))} \widehat{L}(TZ) \right\}^{(8)} \\
&= \frac{8}{3} (4p_1^2 - 7p_2 - D(x)^2).
\end{aligned}$$

We therefore get (3.2), and have completed the proof of Theorem 3.1.

4.3.2. *Proof of Theorem 3.2.* For each i , construct the twisted \widehat{L} -Witten class

$$(4.61) \quad e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \in H^{4*}(Z, \mathbb{Q})$$

and set

$$(4.62) \quad \mathcal{R}_L(P_i, \tau) = \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)}.$$

Lemma 4.4. $\mathcal{R}_L(P_i, \tau)$ is a modular form of weight 10 over $SL(2; \mathbb{Z})$.

Proof. This can be similarly proved as Lemma 4.3 by seeing that

$$\begin{aligned}
& \mathcal{R}_L(P_i, \tau) \\
&= \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \right\}^{(12)} \\
(4.63) \quad &= 2^6 \left\{ e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \left(\prod_{k=1}^6 x_k \frac{\theta'(0, \tau)}{\theta(x_k, \tau)} \frac{\theta_1(x_k, \tau)}{\theta_1(0, \tau)} \frac{\theta_2(x_k, \tau)}{\theta_2(0, \tau)} \frac{\theta_3(x_k, \tau)}{\theta_3(0, \tau)} \right) \right. \\
&\quad \left. \cdot \frac{1}{2} \left(\prod_{l=1}^8 \theta_1(y_l^i, \tau) + \prod_{l=1}^8 \theta_2(y_l^i, \tau) + \prod_{l=1}^8 \theta_3(y_l^i, \tau) \right) \right\}^{(12)},
\end{aligned}$$

and then we can perform the transformation laws (4.6)-(4.9) for the theta functions and the transformation laws (4.21), (4.22) for $E_2(\tau)$ to show that $\mathcal{R}_L(P_i, \tau)$ is a modular form of weight 10 over $SL(2; \mathbb{Z})$. \square

Expanding the q -series, using (4.12), (4.19) and (4.48), we have

$$\begin{aligned}
(4.64) \quad & e^{\frac{1}{24}E_2(\tau)(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(\Phi(T_{\mathbb{C}}Z)) \varphi(\tau)^8 \text{ch}(\mathcal{V}_i) \\
&= \left(e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \right. \\
&\quad \left. - e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \left(-2p_1(TZ) + \frac{1}{30}c_2(W_i) \right) q + O(q^2) \right) \\
&\quad \cdot \widehat{L}(TZ) \text{ch}(\mathbb{C} + D_1q + O(q^2))(1 - 8q + O(q^2))(1 + \text{ch}(W_i)q + O(q^2)) \\
&= e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \\
&\quad + q \left(e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(D_1 - 8 + W_i) - e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \left(-2p_1(TZ) + \frac{1}{30}c_2(W_i) \right) \widehat{L}(TZ) \right) \\
&\quad + O(q^2).
\end{aligned}$$

However modular form of weight 10 must be a multiple of

$$(4.65) \quad E_4(\tau)E_6(\tau) = 1 - 264q + \dots$$

So from (4.64) and (4.65) we have

$$\begin{aligned}
(4.66) \quad & \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \text{ch}(D_1 - 8 + W_i) \right\}^{(12)} \\
&\quad - \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \left(-2p_1(TZ) + \frac{1}{30}c_2(W_i) \right) \widehat{L}(TZ) \right\}^{(12)} \\
&= -264 \left\{ e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \right\}^{(12)}.
\end{aligned}$$

Therefore

$$\begin{aligned}
(4.67) \quad & \left\{ \widehat{L}(TZ) \text{ch}(W_i + D_1 + 256) \right\}^{(12)} \\
& = \left(-2p_1(TZ) + \frac{1}{30}c_2(W_i) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} - 1}{-2p_1(TZ) + \frac{1}{30}c_2(W_i)} \widehat{L}(TZ) \text{ch}(W_i + D_1 + 256) + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \right\}^{(8)}.
\end{aligned}$$

Plugging in D_1 (see (4.57)), we have

$$\begin{aligned}
(4.68) \quad & \left\{ \widehat{L}(TZ) \text{ch}(W_i + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) + 244) \right\}^{(12)} \\
& = \left(-2p_1(TZ) + \frac{1}{30}c_2(W_i) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} - 1}{-2p_1(TZ) + \frac{1}{30}c_2(W_i)} \widehat{L}(TZ) \text{ch}(W_i + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) + 244) \right. \\
& \quad \left. + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(W_i))} \widehat{L}(TZ) \right\}^{(8)}.
\end{aligned}$$

Taking $W_i = V_{\mathbb{C}}(x)$, we have

$$\begin{aligned}
(4.69) \quad & \left\{ \widehat{L}(TZ) \text{ch}(V_{\mathbb{C}}(x) + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) + 244) \right\}^{(12)} \\
& = \left(-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)) \right) \\
& \cdot \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)))} - 1}{-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x))} \widehat{L}(TZ) \text{ch}(\mathfrak{D}) + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)))} \widehat{L}(TZ) \right\}^{(8)},
\end{aligned}$$

where

$$\mathfrak{D} = V_{\mathbb{C}}(x) + 2T_{\mathbb{C}}Z + \wedge^2(T_{\mathbb{C}}Z) - S^2(T_{\mathbb{C}}Z) + 244.$$

Note that the 4-form

$$-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)) = 2\tilde{D}(x).$$

A direct computation shows that the 8-form in the right side of (4.69) verifies

$$\begin{aligned}
& \left\{ -\frac{e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)))} - 1}{-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x))} \widehat{L}(TZ) \text{ch}(\mathfrak{D}) + e^{\frac{1}{24}(-2p_1(TZ) + \frac{1}{30}c_2(V_{\mathbb{C}}(x)))} \widehat{L}(TZ) \right\}^{(8)} \\
& = \frac{8}{3} \left(p_1^2 - 7p_2 - 6p_1\tilde{D}(x) - 4\tilde{D}(x)^2 \right).
\end{aligned}$$

We then get (3.3), and have completed the proof of Theorem 3.2.

5. THE CHARACTERISTIC CLASSES IN THE CUBIC FORMS

The spin^c characteristic classes are determined by Duan [10] by computing the integral cohomology of its classifying space $B\text{Spin}^c$. Let $c \in H^2(B\text{Spin}^c)$ be the class with the mod 2 reduction

$$c \equiv w_2 \pmod{2},$$

where w_2 is the second Stiefel-Whitney class. Then by a theorem of Duan [10] (c.f. Thomas [29])

$$(5.1) \quad H^*(BSpin^c) \cong \mathbb{Z}[c, q_1, q_2, q_3, \dots] \oplus (\text{the 2-torsion part}),$$

where q_i is called the *the i -th universal spin^c class* with $\deg(q_i) = 4i$. The spin^c classes determine the Pontrjagin classes. In the low dimensions, we have

$$(5.2) \quad \begin{aligned} p_1 &= 2q_1 + c^2, \\ p_2 &= 2q_2 + q_1^2, \\ p_3 &= q_3. \end{aligned}$$

The relations between spin^c classes and Stiefel-Whitney classes can be described by the mod 2 reductions of spin^c classes (c.f. Benson-Wood [6]). In the low dimensions,

$$(5.3) \quad \begin{aligned} q_1 &\equiv w_4 \pmod{2}, \\ q_2 &\equiv w_8 \pmod{2}, \\ q_3 &\equiv w_6^2 \pmod{2}. \end{aligned}$$

To obtain spin characteristic classes, we can simply let $c = 0$ in (5.1) and (5.2). In this case, (5.3) is still valid. Notice that in Freed-Hopkins [11], they denoted q_1 by λ and q_2 by p . With the above in hand, it is easy to calculate the following

Theorem 5.1. *Let M be any spin^c manifold with the determinant class $c \in H^2(M)$.*

(i) *One has*

$$4p_2 - p_1^2 - 6p_1c^2 + 39c^4 = 8(q_2 - 2q_1c^2 + 4c^4).$$

Hence,

$$p_c = \frac{4p_2 - p_1^2 - 6p_1c^2 + 39c^4}{8}$$

is well defined and

$$p_c \equiv w_8 \pmod{2}.$$

(ii) *One has*

$$4p_2 - 7p_1^2 + 30p_1c^2 - 15c^4 = 8(q_2 - 3q_1^2 + 4q_1c^2 + c^4).$$

Hence

$$\tilde{p}_c = \frac{4p_2 - 7p_1^2 + 30p_1c^2 - 15c^4}{8}$$

is well defined and

$$\tilde{p}_c \equiv w_8 + w_4^2 + w_2^4 \pmod{2}.$$

(iii) *p_c and \tilde{p}_c satisfy*

$$\tilde{p}_c = p_c - 3\lambda_c^2,$$

where

$$\lambda_c = \frac{1}{2}(p_1 - 3c^2) = q_1 - c^2,$$

and

$$\lambda_c \equiv w_4 + w_2^2 \pmod{2}.$$

Theorem 5.2. *For any orientable manifold,*

$$\begin{aligned} 4p_1^2 - 7p_2 &\equiv w_4^2 \pmod{2}, \\ p_1^2 - 7p_2 &\equiv w_2^4 + w_4^2 \pmod{2}. \end{aligned}$$

6. OBSTRUCTION CLASSES OF spin^ξ AND spin^{w_2} STRUCTURES

Definition 6.1. An oriented closed manifold M is called spin^ξ if its second Stiefel-Whitney class $w_2(M)$ can be realized as that of some real vector bundle ξ of rank 2 over M , that is,

$$w_2(M) = w_2(\xi).$$

The concept spin^ξ is a generalization of spin^c . Indeed, M is spin^c when ξ can be chosen to be orientable. However, there are non- spin^c spin^ξ manifolds.

Definition 6.2. An oriented closed manifold M is called spin^{w_2} if $w_2(M)$ can be realized as that of some nonorientable real vector bundle ξ of rank 2 over M .

Recall that the obstruction class to spin^c structure is the third integral Stiefel-Whitney class $W_3(M) \in H^3(M; \mathbb{Z})$. In contrast, we need to use *cohomology with local coefficients* (or *twisted cohomology*) to investigate the obstructions of spin^ξ and spin^{w_2} structures. There are two standard ways to define cohomology with local coefficients: via module over the group ring of fundamental group, or via *bundle of groups* (for instance, see Hatcher, Section 3.H of [20]). They correspond to each other in a natural way.

Here, we deal with the obstruction problem for the homotopy lifting diagram

$$(6.1) \quad \begin{array}{ccc} & & BO(2) \\ & \nearrow f & \downarrow w_2 \\ M & \xrightarrow{w_2(M)} & K(\mathbb{Z}/2, 2), \end{array}$$

where $K(\mathbb{Z}/2, 2)$ is Eilenberg-MacLane space, and w_2 represents the second universal Stiefel-Whitney class. For the classifying space $BO(2)$, the first universal Stiefel-Whitney class

$$\omega_1 \in H^1(BO(2); \mathbb{Z}/2) \cong \text{Hom}(\pi_1(BO(2)), \text{Aut}(\mathbb{Z}) \cong \mathbb{Z}/2)$$

determines a bundle of groups $\mathbb{Z}^{\omega_1} \rightarrow BO(2)$ with fibre \mathbb{Z} . Moreover, $BO(2)$ is a *generalized Eilenberg-MacLane space* in the sense of Gitler [13], and by Theorem 7.18 of [13], for any $t \in H^1(M; \mathbb{Z}/2) \cong \text{Hom}(\pi_1(M), \pi_1(BO(2)))$

$$(6.2) \quad [M, BO(2)]_t \cong H^2(M; \mathbb{Z}^t),$$

where the set $[M, BO(2)]_t \subseteq [M, BO(2)]$ consists of the classes of maps f such that $f_* = t$ on fundamental groups, and the local coefficient (or the bundle of groups) $\mathbb{Z}^t = t^*(\mathbb{Z}^{\omega_1})$. From the short exact sequence of bundles of groups

$$0 \rightarrow \mathbb{Z}^t \xrightarrow{2} \mathbb{Z}^t \rightarrow \mathbb{Z}/2 \rightarrow 0,$$

there exists a long exact sequence of cohomology with local coefficients

$$(6.3) \quad \cdots \rightarrow H^2(M; \mathbb{Z}^t) \xrightarrow{2} H^2(M; \mathbb{Z}^t) \xrightarrow{\rho_2} H^2(M; \mathbb{Z}/2) \xrightarrow{\beta^t} H^3(M; \mathbb{Z}^t) \rightarrow \cdots,$$

where ρ_2 is the mod-2 reduction to the cohomology with the untwisted coefficient $\mathbb{Z}/2$, and β^t is the t -twisted Bockstein homomorphism. Let us call $W_3^t(M) := \beta^t(w_2(M))$ the *third t -twisted integral Stiefel-Whitney class*. Set $M = BO$ and $t = \omega_1$, we have the *third universal t -twisted integral Stiefel-Whitney class*

$$W_3^{\omega_1} := \beta^{\omega_1}(w_2).$$

In particular, for $t = 0$ $W_3^0(M) = W_3(M)$ is the usual third integral Stiefel-Whitney class.

Theorem 6.1. *An oriented closed manifold M is spin^ξ if and only if*

$$W_3^t(M) = 0,$$

for some $t \in H^1(M; \mathbb{Z}/2)$.

Proof. From the exactness of the sequence (6.3), $W_3^t(M) = 0$ is equivalent to that $w_2(M) = \rho_2(c^t)$ for some $c^t \in H^2(M; \mathbb{Z}^t)$, which by (6.2) is equivalent to the existence of a real vector bundle ξ^t of rank 2 such that $\omega_1(\xi^t) = t$ and $w_2(\xi^t) = w_2(M)$. \square

In particular, the theorem recovers the obstruction result for spin^c structure and determines the obstruction class for spin^{w_2} structure as well.

Corollary 6.1. *An oriented closed manifold M is spin^{w_2} if and only if*

$$W_3^t(M) = 0,$$

for some $t \in H^1(M; \mathbb{Z}/2)$ and $t \neq 0$.

REFERENCES

- [1] L. Alvarez-Gaumé and E. Witten, Gravitational anomalies, *Nucl. Physics* B234 (1983), 269-330.
- [2] M. F. Atiyah, V. K. Patodi and I. M. Singer, Spectral asymmetry and Riemannian geometry I. *Proc. Cambridge Philos. Soc.* 77 (1975), 4369.
- [3] M. F. Atiyah, *K-theory*. Benjamin, New York, 1967.
- [4] M. F. Atiyah and I. M. Singer, The index of elliptic operators III, *Ann of Math.* 87 (1968), 546-604.
- [5] M. F. Atiyah and I. M. Singer, The index of elliptic operators V, *Ann of Math.* 93 (1971), 1391-149.
- [6] D. J. Benson and J. A. Wood, Integral invariants and cohomology of $BSpin(n)$, *Topology* 34 (1995), 13-28.
- [7] J. H. Bruinier, G. van der Geer, G. Harder and D. Zagier, *The 1-2-3 of Modular forms*. Lectures at a Summer School in Nordfjordeid, Norway. Springer, 2007.
- [8] K. Chandrasekharan, *Elliptic Functions*. Springer-Verlag, 1985.
- [9] Q. Chen, F. Han and W. Zhang, Generalized Witten genus and vanishing theorems, *J. Diff. Geom.* 88 (2011), 1-39.
- [10] H. Duan, The characteristic classes and Weyl invariants of Spinor groups, *arXiv:1810.03799*.
- [11] D. S. Freed and M. J. Hopkins, Consistency of M -Theory on nonorientable manifolds, *arXiv:1908.09916*.
- [12] T. Gannon and C. S. Lam, Lattices and θ -function identities. II. Theta series, *J. Math. Phys.* 33 (1992), 871-887.
- [13] S. Gitler, Cohomology operations with local coefficients, *Amer. J. Math.* 85 (1963), 156-188.
- [14] M. B. Green and J. H. Schwarz, Anomaly cancellations in supersymmetric d=10 gauge theory and superstring theory, *Physics Letters* B149 (1984), 117-122.
- [15] V. Gritsenko, Complex vector bundles and Jacobi forms, *Automorphic forms and L-functions (Kyoto, 1999)*. *Srikaisekikenkyusho Kkyroku* No. 1103 (1999), 7185.
- [16] F. Han and K. Liu, Gravitational anomaly cancellation and modular invariance, *Alg. Geom. Top.* 14 (2014), 91-113.
- [17] F. Han, K. Liu and W. Zhang, Modular forms and generalized anomaly cancellation formulas, *J. Geom. Phys.* 62 (2012), 1038-1053.
- [18] F. Han, K. Liu and W. Zhang, Anomaly cancellation and modularity, *Frontiers in Differential Geometry, Partial Differential Equations and Mathematical Physics, In Memory of Gu Chaohao*. pp. 87-104, World Scientific Publishing Co., Inc. 2014.
- [19] F. Han, K. Liu and W. Zhang, Anomaly cancellation and modularity II. The $E_8 \times E_8$ case, *Sci. China Math.* 60 (2017), no. 6, 985994.
- [20] A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002. xii+544 pp.
- [21] M. J. Hopkins, Algebraic topology and modular forms, *Plenary talk, ICM*. Beijing, 2002.
- [22] P. Hořava and E. Witten, Heterotic and Type I string dynamics from eleven dimensions, *Nucl. Phys.* B460 (1996) 506.
- [23] P. Hořava and E. Witten, Eleven dimensional supergravity on a manifold with boundary, *Nucl. Phys.* B475 (1996) 94.

- [24] V. G. Kac, An elucidation of : “Infinite-dimensional algebras, Dedekind’s η -function, classical Moebius function and the very strange formula”. $E_8(1)$ and the cube root of the modular function j , *Adv. in Math.* 35 (1980), 264-273.
- [25] V. G. Kac, *Infinite-Dimensional Lie Algebras*. Cambridge University Press, Cambridge, 3rd edition, 1990.
- [26] K. Liu, Modular invariance and characteristic numbers, *Comm. Math. Phys.* 174 (1995), 29-42.
- [27] K. Liu, On modular invariance and rigidity theorems, *J. Diff. Geom.* 41 (1995), 343-396.
- [28] D. McLaughlin, Orientation and string structures on loop spaces, *Pacific J. Math.* 155 (1992), 143-156.
- [29] E. Thomas, On the cohomology groups of the classifying space for the stable spinor groups, *Bol. Soc. Math. Mex.* 7 (1962), 57-69.
- [30] E. Witten, The index of the Dirac operator in loop space, in *P.S. Landweber, ed., Elliptic Curves and Modular Forms in Algebraic Topology (Proceedings, Princeton 1986) Lecture Notes in Math.* 1326, pp. 161-181, Springer, 1988.
- [31] E. Witten, On flux quantization in M-theory and the effective action, *J. Geom. Phys.* 22 (1997), 1-13.
- [32] D. Zagier, Note on the Landweber-Stong elliptic genus, in *P.S. Landweber, ed., Elliptic Curves and Modular Forms in Algebraic Topology (Proceedings, Princeton 1986), Lecture Notes in Math.* 1326, pp. 216-224, Springer, 1988.
- [33] W. Zhang, Spin^c -manifolds and Rokhlin congruences, *C R Acad Sci Paris, Serie I.* 317 (1993), 689-692.
- [34] W. Zhang, Circle bundles, adiabatic limits of η -invariants and Rokhlin congruences, *Ann Inst Fourier* 44 (1994), 249-270.
- [35] W. Zhang, *Lectures on Chern-Weil Theory and Witten Deformations*. Nankai Tracts in Mathematics Vol. 4, World Scientific, Singapore, 2001.
- [36] W. Zhang, Cobordism and Rokhlin congruences, *Acta Mathematica Scientia* 29B(3) (2009), 609-612.
- [37] W. Zhang, A mod 2 index theorem for pin^- manifolds, *Science China Mathematics* 60 (2017), 1615-1632.

FEI HAN, DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF SINGAPORE, BLOCK S17, 10 LOWER KENT RIDGE ROAD, SINGAPORE 119076 (MATHANF@NUS.EDU.SG)

RUIZHI HUANG, INSTITUTE OF MATHEMATICS AND SYSTEMS SCIENCES, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

E-mail address: huangrz@amss.ac.cn

URL: <https://sites.google.com/site/hrzsea>, <https://hrzsea.github.io/Huang-Ruizhi/>

KEFENG LIU, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT LOS ANGELES, LOS ANGELES, CA 90095, USA (LIU@MATH.UCLA.EDU) AND MATHEMATICAL SCIENCE RESEARCH CENTER, CHONGQING UNIVERSITY OF TECHNOLOGY, CHONGQING, 400054, CHINA

WEIPING ZHANG, CHERN INSTITUTE OF MATHEMATICS & LPMC, NANKAI UNIVERSITY, TIANJIN 300071, CHINA. (WEIPING@NANKAI.EDU.CN)