

9. Moreno, E., and Basler, K. (2004). *dMyc* transforms cells into super-competitors. *Cell* 117, 117–129.
10. Suijkerbuijk, S.J., Kolahgar, G., Kucinski, I., and Piddini, E. (2016). Cell competition drives the growth of intestinal adenomas in *Drosophila*. *Curr. Biol.* 26, 428–438.
11. Eichenlaub, T., Cohen, S.M., and Herranz, H. (2016). Cell competition drives the formation of metastatic tumors in a *Drosophila* model of epithelial tumor formation. *Curr. Biol.* 26, 419–427.
12. Martin, F.A., Perez-Garajo, A., and Morata, G. (2009). Apoptosis in *Drosophila*: compensatory proliferation and undead cells. *Int. J. Dev. Biol.* 53, 1341–1347.
13. Kolahgar, G., Suijkerbuijk, S.J., Kucinski, I., Poirier, E.Z., Mansour, S., Simons, B.D., and Piddini, E. (2015). Cell competition modifies adult stem cell and tissue population dynamics in a JAK-STAT-dependent manner. *Dev. Cell* 34, 297–309.
14. Tavori, Y., and Deng, W.M. (2013). Tissue repair through cell competition and compensatory cellular hypertrophy in postmitotic epithelia. *Dev. Cell* 25, 350–363.
15. Tsuboi, A., Ohsawa, S., Umetsu, D., Sando, Y., Kuranaga, E., Igaki, T., and Fujimoto, K. (2018). Competition for space is controlled by apoptosis-induced change of local epithelial topology. *Curr. Biol.* 28, 2115–2128.
16. Legoff, L., Rouault, H., and Lecuit, T. (2013). A global pattern of mechanical stress polarizes cell divisions and cell shape in the growing *Drosophila* wing disc. *Development* 140, 4051–4059.
17. Mao, Y., Tournier, A.L., Hoppe, A., Kester, L., Thompson, B.J., and Tapon, N. (2013). Differential proliferation rates generate patterns of mechanical tension that orient tissue growth. *EMBO J.* 32, 2790–2803.
18. Shraiman, B.I. (2005). Mechanical feedback as a possible regulator of tissue growth. *Proc. Natl. Acad. Sci. USA* 102, 3318–3323.
19. Curran, S., Strandkvist, C., Bathmann, J., de Gennes, M., Kabla, A., Salbreux, G., and Baum, B. (2017). Myosin II controls junction fluctuations to guide epithelial tissue ordering. *Dev. Cell* 43, 480–492 e486.
20. Aragona, M., Panciera, T., Manfrin, A., Giullitti, S., Michielin, F., Elvassore, N., Dupont, S., and Piccolo, S. (2013). A mechanical checkpoint controls multicellular growth through YAP/TAZ regulation by actin-processing factors. *Cell* 154, 1047–1059.

## Communication: Potassium Channels Define the Dialect

Elena O. Gracheva<sup>1,2,3</sup> and Sviatoslav N. Bagriantsev<sup>1</sup>

<sup>1</sup>Department of Cellular & Molecular Physiology, Yale University School of Medicine, New Haven, CT 06520, USA

<sup>2</sup>Department of Neuroscience, Yale University School of Medicine, New Haven, CT 06520, USA

<sup>3</sup>Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medicine, New Haven, CT 06520, USA

Correspondence: [elena.gracheva@yale.edu](mailto:elena.gracheva@yale.edu) (E.O.G.), [slav.bagriantsev@yale.edu](mailto:slav.bagriantsev@yale.edu) (S.N.B.)

<https://doi.org/10.1016/j.cub.2018.05.048>

**Weakly electric fishes use electric pulses to interact with conspecifics, but the molecular origin of species-specific communication is unknown. A new study shows that some properties of the electric fish ‘language’ are dictated by the activity of a voltage-gated potassium channel in electrocytes.**

Since the late 18th century, decades before Michael Faraday laid the foundation of the theory of electromagnetism, scientists knew that certain species of fish could generate electricity. Some species, such as the electric eel, use strong electric discharges of up to hundreds of volts to stun their prey [1,2]. To produce electricity, the eel uses the electric organ, a unique structure composed of a series of biological batteries. The ‘batteries’, called electrocytes, are modified non-contractile myocytes, the cells that normally build muscles [3]. Similar electric organs were also found in a number of then obscure fish species from the rivers of Africa and South America. But in contrast to the eel, these species failed to produce strong electric output, leaving the researchers

puzzled regarding the function and biological significance of these structures. Some went as far as suggesting that the ‘pseudo-electric organs’ of such species evolved “...*most probably not based upon a natural selective basis*” [4].

The conundrum was resolved in the mid-20th century, when Hans Lissmann obtained a single specimen of *Gymnarcus niloticus*, and described a most peculiar feature of its behavior: the fish was able to avoid obstacles while swimming backwards [5]. Lissmann hypothesized that this was due to the electric organs in the finger-like tail, which he thought the animal used to detect objects. By placing a pair of electrodes connected to an amplifier into the aquarium, he detected a stream of high-frequency (250–300 Hz)

low-voltage (30 mV) electric discharges [5]. Moreover, he noticed that the weakly electric fish not only produced but also sensed electrical pulses produced elsewhere and located their source. This suggested that unlike the eel, who uses electricity as a weapon, *Gymnarcus* uses it in a more subtle way: to navigate in the environment, detect objects and locate prey [6,7].


Lissmann’s works laid the foundation for what is currently part of a flourishing field of bioelectricity and electroreception. The majority of weakly electric fish belong to the Mormyroidae superfamily and includes Gymnarchidae and Mormyridae. Currently, there are over 220 recognized Mormyroidae species, organized into more than 20 genera [8]. All the species from this superfamily can



sense low-voltage discharges produced by their own electric organ as well as those from other individuals. The electric organ discharges (EODs) have species-specific signatures and serve to mediate communication with conspecifics, provide a means for reproductive isolation, and are used for systematics of weakly electric fishes [9–11].

The temporal features of electric organ discharges serve a purpose similar to words and their order in a human language. The principle components of EODs are the frequency, pattern and shape of the waveform. The frequency and pattern of the EODs can vary depending on the social context, whereas the waveform is species-specific and defined by the morphology, innervation and physiology of the electrocytes [12]. This is particularly evident in the two sister taxons of Mormyroidae. *Gymnarchus niloticus*, the single species of the Gymnarchidae family, generates wide wave-like EODs, while most species from the Mormyridae family generate brief multiphasic EODs (Figure 1) [13]. The EODs result from a concerted discharge of action potentials produced by many individual electrocytes. With the advent of the patch-clamp and voltage-clamp techniques, detailed characterization of the ionic currents underlying electrocyte firing became possible. Similar to neuronal and muscle cells, electrocyte discharge is mainly mediated by voltage-gated sodium and potassium channels [14]. Coordinated shifts in the kinetics of sodium and potassium channel activity were proposed to define EOD pulse shape and duration [15]. However, the identity of the channels as well as species-specific signatures of their activity have remained a mystery. The development of high-throughput sequencing and bioinformatics in the last decade started to reveal the molecular origins of the electric organs' discharge at an unprecedented depth [16–19].

A paper by Harold Zakon and colleagues [11] in this issue of *Current Biology* reports a multidisciplinary approach to zoom in on the molecular origin of the wide versus narrow EOD waveform in Gymnarchidae and Mormyridae, respectively. The authors used massively parallel sequencing in combination with differential transcriptome analyses between muscle



**Figure 1. Communication in electric fish.**

The weakly electric fishes communicate with conspecifics via electric discharges with characteristic waveforms: wide for *Gymnarchus* and narrow for *Brienomyrus*. The plasma membrane of the fish electrocytes contains voltage-gated potassium channels  $K_v1.7a$ , which regulate the duration of the waveform. A patch of negatively charged amino acids near the positively charged voltage-sensing domain of *Brienomyrus*  $K_v1.7a$  facilitates channel opening at more negative voltages, shortening the duration of the electric discharge.

and electric organs of *Gymnarchus niloticus* and several mormyrids, including *Brienomyrus brachyistius*, to identify transcripts enriched in the electric organ. They found *kcna7a*, the transcript encoding the voltage-gated potassium channel  $K_v1.7a$ , to be present in large amounts in the electric organ of *Brienomyrus* and other mormyrids. Electrophysiological analysis showed that mormyrid  $K_v1.7a$  has novel properties, different from its orthologues from *Gymnarchus* and other species. In technical terms, mormyrid  $K_v1.7a$  activates at a hyperpolarized membrane potential and has a steep conductance–voltage dependence (Figure 1). This means that the channel activates shortly after the onset of the action potential, narrowing its duration close to the theoretical minimum. Indeed, when properties of the  $K_v1.7a$  orthologues were plugged into a computational

model of electrocyte firing, the results were consistent with the observed differences between the wide and narrow action potentials of *Gymnarchus* and *Brienomyrus*, respectively [11].

What are the molecular origins that dictate the unconventional behavior of mormyrid  $K_v1.7a$ ? The authors noticed that mormyrid  $K_v1.7a$  has a cluster of negatively charged amino acids in the extracellular loop near the voltage-sensing domain (Figure 1). By swapping this region between the  $K_v1.7a$  orthologues, they found that it is sufficient to interconvert properties of  $K_v1.7a$  between the species. While the exact mechanism by which the negatively charged cluster facilitates  $K_v1.7a$  activation remains to be determined, the authors provided compelling evidence supporting the idea that it acts by altering outer membrane surface charges,

which in turn facilitates movement of the voltage-sensing domain in response to small depolarizations [11]. This paradigm is not without a precedent: a similar mechanism was proposed to be at play in the L-type calcium channel found in the electrosensing ampullary organ in little skate [19]. The apparent similarity between the mechanisms potentiating voltage sensitivity of the ion channels from different classes highlights a common pathway targeted in the course of evolution to confer novel properties via a minimal number of amino acid changes.

The new study raises interesting questions. Mormyrids are a very diverse group, and some species have independently evolved wide EOD pulses similar to *Gymnarchus* [20]. Does  $K_v1.7a$  from these species have properties similar to those of *Gymnarchus*, and if so, what additional compensatory mutations are responsible for these changes? It will also be interesting to know whether  $K_v1.7a$  alone is sufficient to fully change the duration of the action potential discharge, or whether other channels play an equally important role. Recent advances in genomic manipulations, such as the CRISPR-Cas9 genome editing technology, should provide an impetus for studies of non-conventional species that possess unique functional specializations. These studies are in need as they lend a novel standpoint to better understand processes that govern speciation and evolution at the molecular level.

#### REFERENCES

- Catania, K. (2014). The shocking predatory strike of the electric eel. *Science* 346, 1231–1234.
- Catania, K.C. (2017). Electrical potential of leaping eels. *Brain Behav. Evol.* 89, 262–273.
- Szabo, T. (1966). The origin of electric organs of *Electrophorus electricus*. *Anat. Rec.* 155, 103–110.
- Dahlgren, U. (1910). The origin of the electricity tissues in fishes. *Amr. Nat.* 44, 193–202.
- Lissmann, H.W. (1951). Continuous electrical signals from the tail of a fish. *Gymnarchus niloticus* Cuv. *Nature* 167, 201–202.
- Lissmann, H.W. (1958). On the function and evolution of electric organs in fish. *J. Exp. Biol.* 35, 156–191.
- Lissmann, H.W., and Machin, K.E. (1958). The mechanism of object location in *Gymnarchus niloticus* and similar fish. *J. Exp. Biol.* 35, 451–486.
- Rich, M., Sullivan, J.P., and Hopkins, C.D. (2017). Rediscovery and description of *Paramormyrops sphekodes* (Sauvage, 1879) and a new cryptic *Paramormyrops* (Mormyridae: Osteoglossiformes) from the Ogooué River of Gabon using morphometrics, DNA sequencing and electrophysiology. *Zool. J. Linnean Soci.* 180, 613–646.
- Hopkins, C.D., and Bass, A.H. (1981). Temporal coding of species recognition signals in an electric fish. *Science* 212, 85–87.
- Gallant, J.R., Arnegard, M.E., Sullivan, J.P., Carlson, B.A., and Hopkins, C.D. (2011). Signal variation and its morphological correlates in *Paramormyrops kingsleyae* provide insight into the evolution of electrogenic signal diversity in mormyrid electric fish. *J. Comp. Physiol. A Neuroethol. Sens. Neural. Behav. Physiol.* 197, 799–817.
- Swapna, I., Ghezzi, A., York, J.M., Markham, M.R., Halling, D.B., Lu, Y., Gallant, J.R., and Zakon, H. (2018). Electrostatic tuning of a potassium channel in electric fish. *Curr. Biol.* 28, 2094–2102.
- Hopkins, C.D. (1995). Convergent designs for electrogenesis and electroreception. *Curr. Opin. Neurobiol.* 5, 769–777.
- Hopkins, C.D. (1999). Design features for electric communication. *J. Exp. Biol.* 202, 1217–1228.
- Shenkel, S., and Sigworth, F.J. (1991). Patch recordings from the electrocytes of *Electrophorus electricus*. Na currents and PNa/PK variability. *J. Gen. Physiol.* 97, 1013–1041.
- Zakon, H., McAnelly, L., Smith, G.T., Dunlap, K., Lopreato, G., Oestreich, J., and Few, W.P. (1999). Plasticity of the electric organ discharge: implications for the regulation of ionic currents. *J. Exp. Biol.* 202, 1409–1416.
- Gallant, J.R., Traeger, L.L., Volkening, J.D., Moffett, H., Chen, P.H., Novina, C.D., Phillips, G.N., Jr., Anand, R., Wells, G.B., Pinch, M., et al. (2014). Nonhuman genetics. Genomic basis for the convergent evolution of electric organs. *Science* 344, 1522–1525.
- Arnegard, M.E., Zwickl, D.J., Lu, Y., and Zakon, H.H. (2010). Old gene duplication facilitates origin and diversification of an innovative communication system—twice. *Proc. Natl. Acad. Sci. USA* 107, 22172–22177.
- Zakon, H.H., Zwickl, D.J., Lu, Y., and Hillis, D.M. (2008). Molecular evolution of communication signals in electric fish. *J. Exp. Biol.* 211, 1814–1818.
- Bellono, N.W., Leitch, D.B., and Julius, D. (2017). Molecular basis of ancestral vertebrate electroreception. *Nature* 543, 391–396.
- Hopkins, C.D. (1981). On the diversity of electric signals in a community of mormyrid electric fish in West Africa. *Am. Zool.* 21, 211–222.

## Social Behavior: How the Brain Thinks like a Mom

Sourish Mukhopadhyay<sup>1,2</sup> and Lisa Stowers<sup>1,\*</sup>

<sup>1</sup>Department of Molecular and Cellular Neuroscience, The Scripps Research Institute, La Jolla, CA, USA

<sup>2</sup>Biomedical Sciences Graduate Program, The Scripps Research Institute, La Jolla, CA, USA

\*Correspondence: [stowers@scripps.edu](mailto:stowers@scripps.edu)

<https://doi.org/10.1016/j.cub.2018.05.049>

Becoming a parent changes our choices and actions. Identifying the underlying neural circuits is necessary to understand the transformation of an animal's behavior post-parenthood. Multiple nodes of the 'parenting circuit' have now been identified to reveal the workings of a single brain region key to the orchestration of parent-specific behaviors.

We begin the first phase of life, through adolescence and beyond selfishly concerned with our own wants and needs; until we become a parent and our behavior radically changes. Caring for

newborns necessitates putting the demands of others ahead of our immediate desires, and requires a change of behavior to prioritize selfless action without immediate personal reward. In

