

A GENERATING FUNCTION FOR BLATTNER'S FORMULA

JEB F. WILLENBRING AND GREGG J. ZUCKERMAN

(Communicated by Gail Letzter)

ABSTRACT. Let G be a connected, semisimple Lie group with finite center and let K be a maximal compact subgroup. We investigate a method to compute multiplicities of K -types in the discrete series using a rational expression for a generating function obtained from Blattner's formula. This expression involves a product with a character of an irreducible finite dimensional representation of K and is valid for any discrete series system. Other results include a new proof of a symmetry of Blattner's formula, and a positivity result for certain low rank examples. We consider in detail the situation for G of type split G_2 . The motivation for this work came from an attempt to understand pictures coming from Blattner's formula, some of which we include in the paper.

1. INTRODUCTION

In [7], a proof of a formula for the restriction of a discrete series representation (see [6]) of a connected, linear, semisimple Lie group to a maximal compact subgroup is given. This formula was first conjectured by Blattner. We recall the formula and its context briefly, from the point of view of root system combinatorics.

Throughout the paper, \mathfrak{g} denotes a semisimple Lie algebra over \mathbb{C} with a fixed Cartan subalgebra \mathfrak{h} . Let $\Phi := \Phi(\mathfrak{g}, \mathfrak{h})$ denote the corresponding root system with Weyl group $W_{\mathfrak{g}}$. Choose a set, Φ^+ , of positive roots and let $\Pi := \{\alpha_1, \dots, \alpha_r\} \subseteq \Phi$ be the simple roots. Let $\Phi^- = -\Phi^+$.

We assume that there exists a function $\theta : \Phi \rightarrow \mathbb{Z}_2$ such that if $\gamma_1, \gamma_2 \in \Phi$ and $\gamma_1 + \gamma_2 \in \Phi$ then $\theta(\gamma_1 + \gamma_2) = \theta(\gamma_1) + \theta(\gamma_2)$. This map provides a \mathbb{Z}_2 -gradation on Φ . We set:

$$\begin{aligned}\Phi_c &:= \{\gamma \in \Phi \mid \theta(\gamma) = 0\}, \\ \Phi_{nc} &:= \{\gamma \in \Phi \mid \theta(\gamma) = 1\}.\end{aligned}$$

Given $\alpha \in \Phi$, set $\mathfrak{g}_{\alpha} = \{X \in \mathfrak{g} \mid [H, X] = \alpha(H)X \ \forall H \in \mathfrak{h}\}$. Let $\mathfrak{k} := \mathfrak{h} \oplus \sum_{\alpha \in \Phi_c} \mathfrak{g}_{\alpha}$ and $\mathfrak{p} := \sum_{\alpha \in \Phi_{nc}} \mathfrak{g}_{\alpha}$. Then, \mathfrak{k} will be a reductive symmetric subalgebra of \mathfrak{g} with $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ the corresponding Cartan decomposition of \mathfrak{g} . As defined, \mathfrak{h} is a Cartan subalgebra for \mathfrak{k} so $\text{rank } \mathfrak{k} = \text{rank } \mathfrak{g}$. Each equal rank symmetric pair corresponds to at least one \mathbb{Z}_2 -gradation in this manner, and conversely.

We shall refer to the elements of Φ_c (resp. Φ_{nc}) as compact (resp. noncompact). The compact roots are a sub-root system of Φ . Let $\Phi_c^+ := \Phi^+ \cap \Phi_c$, $\Phi_{nc}^+ := \Phi^+ \cap \Phi_{nc}$,

Received by the editors April 26, 2007.

2000 *Mathematics Subject Classification.* 22E46 (17B10).

Key words and phrases. Blattner's formula, coherent continuation, discrete series.

The first author was supported in part by NSA Grant # H98230-05-1-0078.

$\Pi_c := \Pi \cap \Phi_c$, and $\Pi_{nc} := \Pi \cap \Phi_{nc}$. Set $\rho_{\mathfrak{g}} := \rho_c + \rho_{nc}$ where $\rho_c := \frac{1}{2} \sum_{\alpha \in \Phi_c^+} \alpha$ and $\rho_{nc} = \frac{1}{2} \sum_{\alpha \in \Phi_{nc}^+} \alpha$. If there is no subscript, we mean $\rho = \rho_c$.

We remark that the \mathbb{Z}_2 -gradation θ is determined by its restriction to Π . Furthermore, to any set partition $\Pi = \Pi_1 \uplus \Pi_2$ there exists a unique \mathbb{Z}_2 -gradation on Φ such that $\Pi_c = \Pi_1$ and $\Pi_{nc} = \Pi_2$.

We denote the Killing form on \mathfrak{g} by $(,)$, which restricts to a nondegenerate form on \mathfrak{h} . Using this form we may define $\iota : \mathfrak{h} \rightarrow \mathfrak{h}^*$ by $\iota(X)(-) = (X, -)$ ($X \in \mathfrak{h}$), which allows us to identify \mathfrak{h} with \mathfrak{h}^* . Under this identification, we have $\iota(H_\alpha) = \frac{2\alpha}{(\alpha, \alpha)} =: \alpha^\vee$, where $H_\alpha \in \mathfrak{h}$ is the simple coroot corresponding to $\alpha \in \Pi$.

For each $\alpha \in \Phi$, set $s_\alpha(\xi) = \xi - (\xi, \alpha^\vee)\alpha$ (for $\xi \in \mathfrak{h}^*$) to be the reflection through the hyperplane defined by α^\vee . For $\alpha_i \in \Pi$, let $s_i := s_{\alpha_i}$, be the simple reflection defined by α_i . Define $\Pi_{\mathfrak{k}}$ to be the set of simple roots in Φ_c^+ and let $W_{\mathfrak{k}}$ denote the Weyl group generated the reflections defined by $\Pi_{\mathfrak{k}}$. Let $W_c = \langle s_\alpha | \alpha \in \Pi_c \rangle$ be the parabolic subgroup of $W_{\mathfrak{g}}$ defined by the compact simple \mathfrak{g} -roots. Note that $W_c \subseteq W_{\mathfrak{k}}$, but we do not have equality in general. For $w \in W_{\mathfrak{k}}$, set $\ell(w) := |w(\Phi_c^+) \cap \Phi_c^-|$. Note that there is also a length function on $W_{\mathfrak{g}}$ (denoted $\ell_{\mathfrak{g}}$) but ℓ refers to $W_{\mathfrak{k}}$.

A weight $\xi \in \mathfrak{h}^*$ is said to be \mathfrak{k} -dominant (resp. \mathfrak{g} -dominant) if $(\xi, \alpha) \geq 0$ for all $\alpha \in \Pi_{\mathfrak{k}}$ (resp. $\alpha \in \Pi$). A weight $\xi \in \mathfrak{h}^*$ is \mathfrak{g} -regular (resp. \mathfrak{k} -regular) if $(\xi, \alpha) \neq 0$ for all $\alpha \in \Phi$ (resp. $\alpha \in \Phi_c$). The integral weight lattice for \mathfrak{g} is denoted by the set $P(\mathfrak{g}) = \{\xi \in \mathfrak{h}^* | (\xi, \alpha^\vee) \in \mathbb{Z} \text{ for all } \alpha \in \Pi_{\mathfrak{g}}\}$. Similarly we let $P(\mathfrak{k})$ denote the abelian group of integral weights for \mathfrak{k} corresponding to $\Pi_{\mathfrak{k}}$. Let the set of \mathfrak{k} - and \mathfrak{g} -dominant integral weights be denoted by $P_+(\mathfrak{k})$ and $P_+(\mathfrak{g})$ respectively. To each element $\delta \in P_+(\mathfrak{k})$ (resp. $P_+(\mathfrak{g})$), let $L_{\mathfrak{k}}(\delta)$ (resp. $L_{\mathfrak{g}}(\delta)$) denote the finite dimensional representation of \mathfrak{k} (resp. \mathfrak{g}) with highest weight δ .

Next, let $Q : P(\mathfrak{k}) \rightarrow \mathbb{N}$ denote the Φ_{nc}^+ -partition function. That is, if $\xi \in \mathfrak{h}^*$ then $Q(\xi)$ is the number of ways of writing ξ as a sum of noncompact positive roots. Put another way: Q defines the coefficients of the product:

$$\sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^\xi = \prod_{\gamma \in \Phi_{nc}^+} (1 - e^\gamma)^{-1}.$$

Finally, we define the Blattner formula. For $\delta, \mu \in P(\mathfrak{k})$,

$$B(\delta, \mu) := \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} Q(w(\delta + \rho) - \rho - \mu).$$

It is convenient to introduce the notation $w.\xi = w(\xi + \rho) - \rho$ for $w \in W_{\mathfrak{k}}$ and $\xi \in \mathfrak{h}^*$. It is easy to see that $B(v.\delta, \mu) = (-1)^{\ell(v)} B(\delta, \mu)$. In light of this fact, we assume that $\delta \in P_+(\mathfrak{k})$.

Historically, Blattner's formula arises out of the study of the discrete series and its generalizations (see [1–3, 5–7, 10–13]).

Theorem 1.1 (See [7]). *Assume $\lambda = \lambda(\mu) := \mu - \rho_{nc} + \rho_c$ is \mathfrak{g} -dominant and \mathfrak{g} -regular. Then, $B(\delta, \mu)$ is the multiplicity of the finite dimensional \mathfrak{k} -representation, $L_{\mathfrak{k}}(\delta)$, in the discrete series representation of G with Harish-Chandra parameter λ .*

In this paper, we do not impose the \mathfrak{g} -dominant regular condition on $\lambda(\mu)$. This is natural from the point of view of representation theory as it is related to the coherent continuation of the discrete series (see [10], [11] and [14]).

From our point of view, the goal is to understand the Blattner formula in as combinatorial fashion as possible. Thus it is convenient to introduce the following generating function:

Definition: For $\delta \in P_+(\mathfrak{k})$ we define the formal series:

$$\mathbf{b}(\delta) := \sum_{\mu \in \mathfrak{h}^*} B(\delta, \mu) e^\mu.$$

The main result of this paper is Proposition 2.1 of Section 2, which states: For $\delta \in P_+(\mathfrak{k})$,

$$\mathbf{b}(\delta) = \text{ch}L_{\mathfrak{k}}(\delta) \frac{\prod_{\gamma \in \Phi_c^+} (1 - e^{-\gamma})}{\prod_{\gamma \in \Phi_{nc}^+} (1 - e^{-\gamma})},$$

where $\text{ch}L_{\mathfrak{k}}(\delta)$ denotes the character of $L_{\mathfrak{k}}(\delta)$.

Of particular interest are the cases where $\Pi_c \neq \emptyset$, which we address in Section 3. From the point of view of representation theory these include, for example, the holomorphic and Borel-de Siebenthal discrete series (see [7]). More recently, the latter has been addressed in [5].

The Blattner formula for the case of $\Pi_c = \emptyset$ is often particularly difficult to compute explicitly when compared to, say, the cases corresponding to holomorphic discrete series. The $\Pi_c = \emptyset$ case corresponds to the *generic* discrete series of the corresponding real semisimple Lie group. In Section 4 we explore this situation in some detail for the Lie algebra G_2 .

Finally, in light of Theorem 1.1 one may observe that if $\delta \in P_+(\mathfrak{k})$ and $\lambda(\mu)$ is \mathfrak{g} -dominant regular then $B(\delta, \mu) \geq 0$. Our goal is to investigate the positivity of Blattner's formula using combinatorial methods. Of particular interest is the positivity when we relax the \mathfrak{g} -dominance condition on $\lambda(\mu)$. Some results in this direction are suggested by the recent work of Penkov and the second author (see [9]).

In Section 3 we prove the existence of a skew symmetry of Blattner's formula that exists whenever $\Pi_c \neq \emptyset$. Thus, the condition that $\Pi_c = \emptyset$ is necessary for $B(\delta, \mu) \geq 0$ for all $\delta \in P_+(\mathfrak{k})$ and $\mu \in P(\mathfrak{k})$. In the situation where $\Pi_c = \emptyset$ we introduce the following:

Definition: We say that a semisimple Lie algebra is **b**-positive if the Blattner formula corresponding to the \mathbb{Z}_2 -gradation with $\Pi_c = \emptyset$ has the property that:

$$B(\delta, \mu) \geq 0 \quad \text{for all } \delta \in P_+(\mathfrak{k}) \text{ and } \mu \in P(\mathfrak{k}).$$

The terminology stems from the fact that a simple Lie algebra is **b**-positive if and only if the coefficients of $\mathbf{b}(\delta)$ are non-negative for all $\delta \in P_+(\mathfrak{k})$. Since the character of $L_{\mathfrak{k}}(\delta)$ can be written as a non-negative integer combination of characters of T , we have that $\mathbf{b}(\delta)$ has non-negative integer coefficients if $\mathbf{b}(0)$ does. Thus the question of **b**-positivity reduces to the case of $\delta = 0$. In Section 5, it is shown that the only **b**-positive simple Lie algebras are of type A_1 , A_2 , A_3 , B_2 , C_3 , D_4 and G_2 . We prove this result by examining the coefficients of $\mathbf{b}(0)$.

2. PROOF OF THE MAIN RESULT

Proposition 2.1. For $\delta \in P_+(\mathfrak{k})$,

$$\mathbf{b}(\delta) = \text{ch } L_{\mathfrak{k}}(\delta) \frac{\prod_{\gamma \in \Phi_c^+} (1 - e^{-\gamma})}{\prod_{\gamma \in \Phi_{nc}^+} (1 - e^{-\gamma})},$$

where $\text{ch } L_{\mathfrak{k}}(\delta)$ denotes the character of $L_{\mathfrak{k}}(\delta)$.

Proof. From the definition of Blattner's formula we have:

$$\mathbf{b}(\delta) = \sum_{\mu \in \mathfrak{h}^*} \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} Q(w(\delta + \rho) - \rho - \mu) e^{\mu}.$$

First we make the substitution, $\mu = w(\delta + \rho) - \rho - \xi$, and reorganize the sum:

$$\begin{aligned} \mathbf{b}(\delta) &= \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} \sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^{w(\delta + \rho) - \rho - \xi} \\ &= \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} e^{w(\delta + \rho) - \rho} \sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^{-\xi} \\ &= \frac{\sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} e^{w(\delta + \rho) - \rho}}{\prod_{\gamma \in \Phi_{nc}^+} (1 - e^{-\gamma})} \\ &= \frac{\sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} e^{w(\delta + \rho) - \rho} \prod_{\gamma \in \Phi_c^+} (1 - e^{-\gamma})}{\prod_{\gamma \in \Phi_c^+} (1 - e^{-\gamma}) \prod_{\gamma \in \Phi_{nc}^+} (1 - e^{-\gamma})}. \end{aligned}$$

As is well known, the character may be expressed using Weyl's formula (see [4, 8]) as in the following:

$$\text{ch } L_{\mathfrak{k}}(\delta) = \frac{\sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} e^{w(\delta + \rho) - \rho}}{\prod_{\gamma \in \Phi_c^+} (1 - e^{-\gamma})}.$$

□

The result allows us to compute the Blattner formula as follows:

$$\sum_{\mu \in \mathfrak{h}^*} B(\delta, \mu) e^{\mu} = \text{ch } L_{\mathfrak{k}}(\delta) \sum_{\nu \in \mathfrak{h}^*} B(0, \nu) e^{\nu}.$$

Let $\text{ch } L_{\mathfrak{k}}(\delta) = \sum_{\gamma \in \mathfrak{h}^*} m_{\gamma} e^{\gamma}$ and we obtain:

$$\sum_{\mu \in \mathfrak{h}^*} B(\delta, \mu) e^{\mu} = \sum_{\gamma, \nu \in \mathfrak{h}^*} m_{\gamma} B(0, \nu) e^{\gamma + \nu}.$$

Thus for $\delta \in P_+(\mathfrak{k})$ and $\mu \in P(\mathfrak{k})$ we have

$$B(\delta, \mu) = \sum_{\gamma \in P(\mathfrak{k})} m_{\gamma} B(0, \mu - \gamma).$$

Note that the numbers m_{γ} are weight multiplicities for the representation $L_{\mathfrak{k}}(\delta)$.

3. (SKEW-)SYMMETRIES OF BLATTNER'S FORMULA

The main result of this section is:

Proposition 3.1. For $v \in W_c$:

$$\begin{aligned} B(\delta, \mu) &= B(\delta, v \cdot \mu) && \text{if } \ell(v) \text{ is even,} \\ B(\delta, \mu) &= -B(\delta, v \cdot \mu) && \text{if } \ell(v) \text{ is odd.} \end{aligned}$$

Although this is well known to experts, we include our proof as it requires very little technical machinery.

Definition: For $w \in W_{\mathfrak{g}}$ and $\xi \in \mathfrak{h}^*$ let $Q_w(\xi) := Q(w^{-1}\xi)$.

Lemma 3.1. *If $w(\Phi_{nc}^+) = \Phi_{nc}^+$ then $Q_w = Q$.*

Proof. It is enough to show

$$\sum_{\xi \in \mathfrak{h}^*} Q_w(\xi) e^{\xi} = \sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^{\xi},$$

which follows from the following calculation:

$$\begin{aligned} \sum_{\xi} Q_w(\xi) e^{\xi} &= \sum_{\xi} Q(\xi) e^{w(\xi)} &&= w \left(\sum_{\xi} Q(\xi) e^{\xi} \right) \\ &= w \left(\prod_{\alpha \in \Phi_{nc}^+} (1 - e^{\alpha})^{-1} \right) &&= \prod_{\alpha \in \Phi_{nc}^+} (1 - e^{w(\alpha)})^{-1} \\ &= \prod_{w^{-1}(\alpha) \in \Phi_{nc}^+} (1 - e^{\alpha})^{-1} &&= \prod_{\alpha \in w(\Phi_{nc}^+)} (1 - e^{\alpha})^{-1}. \end{aligned}$$

□

Lemma 3.2. *For all $w \in W_c$, $w(\Phi_{nc}^+) = \Phi_{nc}^+$.*

Proof. Note that W_c is the Weyl group of a reductive Levi factor \mathfrak{l} of a parabolic subalgebra $\mathfrak{q} \subseteq \mathfrak{g}$. We have a generalized triangular decomposition $\mathfrak{g} = \mathfrak{u}^- \oplus \mathfrak{l} \oplus \mathfrak{u}^+$ with $\mathfrak{q} = \mathfrak{l} \oplus \mathfrak{u}^+$. The noncompact root spaces contained in \mathfrak{u}^+ are positive. Furthermore, all noncompact positive roots spaces are contained in \mathfrak{u}^+ because $\mathfrak{l} \subseteq \mathfrak{k}$. The Lie algebra \mathfrak{u}^+ is an \mathfrak{l} -module and therefore the weights are preserved by W_c . It is this fact that implies that W_c takes positive noncompact roots to positive roots. We now need to show that W_c takes noncompact roots to noncompact roots.

For roots $\beta \in \Phi_{nc}$ and $\alpha \in \Phi_c$, we have the formula $s_{\alpha}(\beta) = \beta - \frac{2(\alpha, \beta)}{(\alpha, \alpha)} \alpha$ with $\frac{2(\alpha, \beta)}{(\alpha, \alpha)}$ an integer. Thus, the reflection of a noncompact root across a hyperplane defined by a compact root is noncompact. The reflections generate W_c . □

Proof of Proposition 3.1. Let $v \in W_c$:

$$B(\delta, \mu) = \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} Q(w \cdot \delta - \mu) = \sum_{w \in v(W_{\mathfrak{k}})} (-1)^{\ell(v^{-1}w)} Q((v^{-1}w) \cdot \delta - \mu).$$

By definition, $\ell(v) = \ell(v^{-1})$. Combining this with the definition of the “dot” action we obtain:

$$\begin{aligned} B(\delta, \mu) &= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} Q(v^{-1}w(\delta + \rho) - \rho - \mu) \\ &= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{k}}} (-1)^{\ell(w)} Q(v(v^{-1}w(\delta + \rho) - \rho - \mu)). \end{aligned}$$

Next, we use the fact that $Q = Q_{v^{-1}}$ using Lemmas 3.1 and 3.2. The rest is a calculation.

$$\begin{aligned}
B(\delta, \mu) &= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{t}}} (-1)^{\ell(w)} Q(w(\delta + \rho) - v(\mu + \rho)) \\
&= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{t}}} (-1)^{\ell(w)} Q(w(\delta + \rho) - \rho - v(\mu + \rho) + \rho) \\
&= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{t}}} (-1)^{\ell(w)} Q(w(\delta + \rho) - \rho - (v(\mu + \rho) - \rho)) \\
&= (-1)^{\ell(v)} \sum_{w \in W_{\mathfrak{t}}} (-1)^{\ell(w)} Q(w.\delta - v.\mu) = (-1)^{\ell(v)} B(\delta, v.\mu).
\end{aligned}$$

□

4. THE CASE OF G_2

The following is a complete calculation of $\mathbf{b}(0)$ for the case of the Lie algebra $\mathfrak{g} := G_2$ when $\Pi_c = \emptyset$. Let α and β be a choice of noncompact simple roots for G_2 with α long and β short. The compact positive roots are $\Phi_c^+ = \{\alpha + \beta, \alpha + 3\beta\}$, while the noncompact positive roots are $\Phi_{nc}^+ = \{\alpha, \beta, \alpha + 2\beta, 2\alpha + 3\beta\}$. Denote the Φ_{nc}^+ -partition function by $Q : \mathfrak{h}^* \rightarrow \mathbb{Z}$. We have:

$$\mathbf{q} := \sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^{-\xi} = \prod_{\gamma \in \Phi_{nc}^+} (1 - e^{-\gamma})^{-1}.$$

Let $x = e^{-\alpha}$ and $y = e^{-\beta}$. Thus,

$$\mathbf{q} = \frac{1}{(1-x)(1-y)(1-xy^2)(1-x^2y^3)}.$$

Let the simple reflection corresponding to $\alpha + \beta$ (resp. $\alpha + 3\beta$) be s_1 (resp. s_2). We have four terms in Blattner's formula for $\delta = 0$:

$$B(0, \mu) = Q(-\mu) - Q(s_1\rho - \rho - \mu) - Q(s_2\rho - \rho - \mu) + Q(s_1s_2\rho - \rho - \mu),$$

and our goal will be to close the sum $\mathbf{b}(0) := \sum_{\mu \in \mathfrak{h}^*} B(0, \mu) e^{\mu}$. We will do this by multiplying by e^{μ} and summing over μ for each of the four terms. Observe that:

$$\sum_{\xi \in \mathfrak{h}^*} Q(\xi) e^{-\xi} = \sum_{\xi \in \mathfrak{h}^*} Q(-\xi) e^{\xi}.$$

Thus, $\mathbf{q} = \sum_{\xi \in \mathfrak{h}^*} Q(-\xi) e^{\xi}$. Next consider the sum:

$$T_1 := \sum_{\xi \in \mathfrak{h}^*} Q(s_1\rho - \rho - \xi) e^{\xi}.$$

We make the substitution $-\mu = s_1\rho - \rho - \xi$ so that the above sum becomes:

$$\sum_{\mu \in \mathfrak{h}^*} Q(-\mu) e^{s_1\rho - \rho + \mu} = e^{s_1\rho - \rho} \sum_{\mu \in \mathfrak{h}^*} Q(-\mu) e^{\mu}.$$

Thus the above sum is equal to $e^{s_1\rho - \rho} \mathbf{q}$, which we denote by \mathbf{T}_1 . Similarly, we set $\mathbf{T}_2 := e^{s_2\rho - \rho} \mathbf{q}$ and $\mathbf{T}_3 := e^{s_1s_2\rho - \rho} \mathbf{q}$, and $\mathbf{T}_0 := \mathbf{q}$. Thus, $\mathbf{b}(0) = \mathbf{T}_0 - \mathbf{T}_1 - \mathbf{T}_2 + \mathbf{T}_3$.

Now we write the above in terms of x and y . Note that $\rho = \alpha + 2\beta$, $s_1\rho = \beta$, $s_2\rho = -\beta$, and from these we can easily see:

$$e^{s_1\rho - \rho} = xy, \quad e^{s_2\rho - \rho} = xy^3, \quad \text{and} \quad e^{s_1s_2\rho - \rho} = x^2y^4.$$

Putting everything together we see $\mathbf{b}(0) = (1 - xy)(1 - xy^3)\mathbf{q}$. Or equivalently, if $\mu = -k\alpha - \ell\beta$ then the value of $B(0, \mu)$ is the coefficient of $x^k y^\ell$ in:

$$\mathbf{b}(0) = \frac{(1 - xy)(1 - xy^3)}{(1 - x)(1 - y)(1 - xy^2)(1 - x^2y^3)} = \frac{1}{(1 - x)(1 - xy^2)} + \frac{y}{(1 - y)(1 - x^2y^3)}.$$

Note that in the latter expression, it is clear that the coefficients in the series are positive. The positivity of the coefficients of $\mathbf{b}(\delta)$ follows from the positivity of the coefficients of $\mathbf{b}(0)$. The question of positivity for a general semisimple Lie algebra will be addressed in Section 5.

It is important to note that as we change Π_c , the value of $\mathbf{b}(0)$ changes as well. For example, when $\Pi_c = \{\beta\}$,

$$\mathbf{b}(0) = \frac{(1 - y)(1 - x^2y^3)}{(1 - x)(1 - xy)(1 - xy^2)(1 - xy^3)} = \frac{1}{(1 - x)(1 - xy^2)} - \frac{y}{(1 - xy)(1 - y^3x)},$$

which we can easily see does *not* have non-negative integer coefficients. We leave it as an exercise to the reader to compute $\mathbf{b}(0)$ when $\Pi_c = \{\alpha\}$.

The generating functions for other δ involve multiplying the above product by a polynomial in $x^{\pm\frac{1}{2}}, y^{\pm\frac{1}{2}}$ that represents the character of the corresponding irreducible finite dimensional representation of $\mathfrak{k} = \mathfrak{so}_4$.

One can plot the coefficients of this formal series, as we do next. In all pictures we have labeled the scale on the axes and normalized the short root (β) to have length 1 and be positioned at 3 o'clock. The long root α is at 10 o'clock.

The first two images (Figure 1) are for the case with $\Pi_c = \emptyset$ (generic) and correspond to $\delta = 0$ and δ the highest weight of the standard 4-dimensional representation of \mathfrak{so}_4 . We also display the same two δ 's in the case when $\Pi_c = \{\beta\}$ (Borel-de Siebenthal) in Figures 2. These latter two figures clearly display a skew-symmetry addressed in Section 3.




Figure 1: $\mathbf{b}(\delta)$ when $\Pi_c = \emptyset$ with $\delta = 0$ (left) and $\delta = \rho$ (right)




Figure 2: $\mathbf{b}(\delta)$ when $\Pi_c = \{\beta\}$ with $\delta = 0$ (left) and $\delta = \rho$ (right)

5. \mathbf{b} -POSITIVE SIMPLE LIE ALGEBRAS

Proposition 5.1. *The only \mathbf{b} -positive simple Lie algebras are of type $A_1, A_2, A_3, B_2, C_3, D_4$ and G_2 .*

To prove the proposition we use the following lemma to exclude the other cases.

Lemma 5.1. *Let \mathfrak{l} be the semisimple Levi factor of a parabolic subalgebra of \mathfrak{g} . If \mathfrak{g} is \mathbf{b} -positive then \mathfrak{l} is also \mathbf{b} -positive.*

Proof. It is a consequence of the main proposition that for any rank r simple Lie algebra \mathfrak{g} we may express $\mathbf{b}(0)$ as a rational function in x_1, \dots, x_r where $x_i = e^{-\alpha_i}$ (recall that α_i is the i^{th} -simple root for \mathfrak{g}). For any j , if we set $x_j = 0$ in $\mathbf{b}(0)$ then the resulting expression is $\mathbf{b}(0)$ for a semisimple subalgebra of \mathfrak{g} . It is not hard to see that this subalgebra is the Levi factor of the maximal parabolic of \mathfrak{g} corresponding to α_j . More generally, let $S \subseteq \Pi$. If we set $x_i = 0$ in $\mathbf{b}(0)$ for all $\alpha_i \in S$ then the resulting expression is $\mathbf{b}(0)$ for the Levi subalgebra, \mathfrak{l}_S , of the corresponding parabolic. Note that the terms in the series expansion for $\mathbf{b}(0)$ for \mathfrak{l}_S are also terms in the series expansion of $\mathbf{b}(0)$ for \mathfrak{g} . Thus a negative coefficient in the former implies a negative coefficient in the latter. \square

The proof of Proposition 5.1 involves a case-by-case analysis using the main proposition expressing $\mathbf{b}(0)$ as a product, Lemma 5.1 and the computer algebra package, MAPLE. Recall that the \mathbf{b} -positivity of G_2 was proved in Section 4.

5.1. **Type A.** For \mathfrak{sl}_2 , we have: $\mathbf{b}(0) = \frac{1}{1-x_1}$. For \mathfrak{sl}_3 :

$$\mathbf{b}(0) = \frac{1 - x_1 x_2}{(1 - x_1)(1 - x_2)} = 1 + \frac{x_1}{1 - x_1} + \frac{x_2}{1 - x_2}.$$

And for \mathfrak{sl}_4 :

$$\begin{aligned} \mathbf{b}(0) &= \frac{(1 - x_1 x_2)(1 - x_2 x_3)}{(1 - x_1)(1 - x_2)(1 - x_3)(1 - x_1 x_2 x_3)} \\ &= \frac{1}{(1 - x_3)(1 - x_1)} + \frac{x_2}{(1 - x_2)(1 - x_1 x_2 x_3)}. \end{aligned}$$

The partial fraction on the right of each of these examples establishes that the coefficients are indeed positive. Next we consider \mathfrak{sl}_5 where we have a negative result. First we note:

$$\mathbf{b}(0) = \frac{(1-x_1x_2)(1-x_2x_3)(1-x_3x_4)(1-x_1x_2x_3x_4)}{(1-x_1)(1-x_2)(1-x_3)(1-x_4)(1-x_1x_2x_3)(1-x_2x_3x_4)},$$

for \mathfrak{sl}_5 . We expand in a formal power series and observe that the coefficient of $x_1x_2^2x_3^2x_4$ is -1 . This means that \mathbf{b} -positivity fails for this example. We then also see failure of \mathbf{b} -positivity for any simple Lie algebra which has A_4 as a Levi factor of a parabolic. Thus, we exclude all higher rank type A examples as well as B_n ($n \geq 5$), C_n ($n \geq 5$), D_n ($n \geq 5$), E_6 , E_7 and E_8 .

5.2. Type B. We only need to examine \mathfrak{so}_5 and \mathfrak{so}_7 . For \mathfrak{so}_5 we have:

$$\begin{aligned} \mathbf{b}(0) &= \frac{1-x_1x_2}{(1-x_1)(1-x_2)(1-x_1x_2^2)} \\ &= \frac{1}{(1-x_2^2)(1-x_1)} + \frac{x_2}{(1-x_2^2)(1-x_1x_2^2)}. \end{aligned}$$

Thus the coefficients of the series expansion are non-negative.

Now consider \mathfrak{so}_7 :

$$\mathbf{b}(0) = \frac{(1-x_1x_2)(1-x_2x_3)(1-x_1x_2x_3^2)}{(1-x_1)(1-x_2)(1-x_3)(1-x_1x_2x_3)(1-x_2x_3^2)(1-x_1x_2^2x_3^2)}.$$

Upon expansion we see that the coefficient of $x_1^2x_2^3x_3^3$ is -1 . Thus, we may exclude this and higher rank type B examples as they have B_3 as the Levi factor of a parabolic. In particular, we may exclude B_4 , which is the only type B example that has not been excluded yet. We also may exclude F_4 for the same reason.

5.3. Type C. We must examine $\mathbf{b}(0)$ for \mathfrak{sp}_6 and \mathfrak{sp}_8 as these are the only examples not yet addressed. For \mathfrak{sp}_6 (ie: C_3), we have:

$$\mathbf{b}(0) = \frac{(1-x_1x_2)(1-x_2x_3)(1-x_1x_2^2x_3)}{(1-x_1)(1-x_2)(1-x_3)(1-x_1x_2x_3)(1-x_2^2x_3)(1-x_1^2x_2^2x_3)}.$$

The coefficients are positive as the above expression is equal to:

$$\begin{aligned} &\frac{1}{(1-x_1)(1-x_2^2x_3)} + \frac{x_3}{(1-x_2^2x_3)(1-x_1^2x_2^2x_3)(1-x_3)} + \frac{x_2}{(1-x_2^2x_3)(1-x_1^2x_2^2x_3)(1-x_2)} \\ &+ \frac{x_1x_3}{(1-x_1)(1-x_3)(1-x_1^2x_2^2x_3)} + \frac{x_1^2x_2^3x_3^2}{(1-x_2^2x_3)(1-x_1^2x_2^2x_3)(1-x_1x_2x_3)}. \end{aligned}$$

However, we do not have \mathbf{b} -positivity for \mathfrak{sp}_8 , as we see -1 as the coefficient of $x_1x_2^3x_3^3x_4^2$ in $\mathbf{b}(0) = \frac{NUM}{DEN}$ where:

$$\begin{aligned} NUM &:= (1-x_1x_2)(1-x_2x_3)(1-x_3x_4)(1-x_1x_2x_3x_4) \\ &\quad (1-x_2x_3^2x_4)(1-x_1x_2^2x_3^2x_4), \\ DEN &:= (1-x_1)(1-x_2)(1-x_3)(1-x_4)(1-x_1x_2x_3)(1-x_2x_3x_4) \\ &\quad (1-x_3^2x_4)(1-x_1x_2x_3^2x_4)(1-x_2^2x_3^2x_4)(1-x_1^2x_2^2x_3^2x_4). \end{aligned}$$

5.4. **Type D.** The only case left is \mathfrak{so}_8 , where we have $\mathbf{b}(0) = \frac{NUM}{DEN}$ where

$$\begin{aligned} NUM &= (1 - x_1 x_2)(1 - x_2 x_3)(1 - x_2 x_4)(1 - x_1 x_2 x_3 x_4), \\ DEN &= (1 - x_1)(1 - x_2)(1 - x_3)(1 - x_4)(1 - x_1 x_2 x_3)(1 - x_1 x_2 x_4) \\ &\quad (1 - x_2 x_3 x_4)(1 - x_1 x_2^2 x_3 x_4). \end{aligned}$$

We see that

$$\mathbf{b}(0) = \frac{1}{\frac{(1 - x_1 x_2^2 x_3 x_4)(1 - x_1)(1 - x_3)(1 - x_4)}{x_2} + \frac{(1 - x_1 x_2 x_4)(1 - x_2)(1 - x_2 x_3 x_4)(1 - x_1 x_2 x_3)}{x_2}}.$$

Thus the coefficients of the series expansion are non-negative integers.

REFERENCES

- [1] Thomas J. Enright, *On the fundamental series of a real semisimple Lie algebra: their irreducibility, resolutions and multiplicity formulae*, Ann. of Math. (2) **110** (1979), no. 1, 1–82.
- [2] Thomas J. Enright and V. S. Varadarajan, *On an infinitesimal characterization of the discrete series*, Ann. of Math. (2) **102** (1975), no. 1, 1–15.
- [3] Thomas J. Enright and Nolan R. Wallach, *The fundamental series of representations of a real semisimple Lie algebra*, Acta Math. **140** (1978), no. 1-2, 1–32.
- [4] R. Goodman and N.R. Wallach, *Representations and invariants of the classical groups*, Cambridge University Press, Cambridge, 1998.
- [5] B. Gross and N. Wallach, *Restriction of small discrete series representations to symmetric subgroups*, The mathematical legacy of Harish-Chandra (Baltimore, MD, 1998), Proc. Sympos. Pure Math., vol. 68, Amer. Math. Soc., Providence, RI, 2000, pp. 255–272.
- [6] Harish-Chandra, *Discrete series for semisimple Lie groups. II. Explicit determination of the characters*, Acta Math. **116** (1966), 1–111.
- [7] Henryk Hecht and Wilfried Schmid, *A proof of Blattner’s conjecture*, Invent. Math. **31** (1975), no. 2, 129–154.
- [8] Anthony W. Knap, *Lie groups beyond an introduction*, Second, Progress in Mathematics, vol. 140, Birkhäuser Boston Inc., Boston, MA, 2002.
- [9] Ivan Penkov and Gregg Zuckerman, *Generalized Harish-Chandra modules with generic minimal \mathfrak{k} -type*, Asian J. Math. **8** (2004), no. 4, 795–811.
- [10] Wilfried Schmid, *L^2 -cohomology and the discrete series*, Ann. of Math. (2) **103** (1976), no. 2, 375–394.
- [11] David A. Vogan Jr., *Representations of real reductive Lie groups*, Progress in Mathematics, vol. 15, Birkhäuser Boston, Mass., 1981.
- [12] Nolan R. Wallach, *Real reductive groups. I*, Pure and Applied Mathematics, vol. 132, Academic Press Inc., Boston, MA, 1988.
- [13] ———, *Real reductive groups. II*, Pure and Applied Mathematics, vol. 132, Academic Press Inc., Boston, MA, 1992.
- [14] Gregg J. Zuckerman, *Coherent translation of characters of semisimple Lie groups*, Proceedings of the International Congress of Mathematicians (Helsinki, 1978), Acad. Sci. Fennica, Helsinki, 1980, pp. 721–724.

JEB F. WILLENBRING, UNIVERSITY OF WISCONSIN - MILWAUKEE; DEPARTMENT OF MATHEMATICAL SCIENCES; P.O. BOX 0413; MILWAUKEE WI 53201-0413

E-mail address: jw@uwm.edu

GREGG J. ZUCKERMAN, YALE UNIVERSITY MATHEMATICS DEPT.; PO Box 208283; NEW HAVEN, CT 06520-8283

E-mail address: gregg.zuckerman@yale.edu