



Rehab Practice Guidelines for: Repair of Large Rotator Cuff Tears

Assumptions:

1. Tears > 2.5 centimeters
2. No retraction
3. Arthroscopic assisted mini-open procedure

Primary Surgery:

- Repair of the supraspinatus & infraspinatus tendons
- Subacromial decompression

Date of Surgery: _____

Secondary Surgery (possible): Distal clavicle excision

Precautions:

- For all passive IR/ER guidelines check with surgeon
- Primary repair (supraspinatus & infraspinatus) - **NO BEHIND THE BACK INTERNAL ROTATION (TOWEL STRETCH)**
- Primary repair – (subacromial decompression) - **No heavy resisted flexion** for 6 weeks
- Primary repair (supraspinatus & split deltoid) - **No resisted abduction** for 8 weeks
- Primary repair (supraspinatus & infraspinatus) - **No resisted external rotation** for 8 weeks
- No additional precautions for distal clavicle excision

Expected # of visits: 19-36

Week Numbers	Treatment Strategies	Milestones ²
<p><u>Week 1-2</u> Dates: _____ to _____</p> <ul style="list-style-type: none"> • No formal PT • Use sling with ABD pillow 24 hrs/day² • No Driving 	<ul style="list-style-type: none"> • Ice for pain and inflammation control • Remove Sling TID for Pendulum exercises (Codman's)² 	<ul style="list-style-type: none"> • Sleep comfortably through the night wearing sling
<p><u>Weeks 3-5</u> Dates: _____ to _____</p> <p>Total Visits: 6-9</p> <ul style="list-style-type: none"> • Begin PT 2-3 visits/week • Continue sling use 24 hrs/day² • D/C ABD pillow 	<ul style="list-style-type: none"> • <u>Modalities for pain and inflammation control as needed</u>³ • <u>Scar Mobilization</u> when incisions are healed • <u>Joint Mobilization</u> <ul style="list-style-type: none"> ○ If hypomobile - grade III/IV mobilizations ○ If normal- grade I/II mobilizations PRN • <u>PROM, AAROM</u> exercises in all planes only to restrictions stated in milestones; no IR in 0° ABD.^{3,6} • <u>Scapular control exercises</u>^{3,4,6} <ul style="list-style-type: none"> ○ Scap PNF ○ T-Band Rows, prone extension to plane of body • <u>Rhythmic stabilization exercises</u>^{2,3,7} • <u>Initiate HEP</u> 	<p><u>PROM:</u></p> <ul style="list-style-type: none"> • ER/IR (90° ABD), HOR ADD: determined by the surgeon • Flexion, ABD: to tolerance up to 90° - 120° • Normal glenohumeral jt. mobility • Normal scapulohumeral rhythm • No hypomobility or hypersensitivity of the scars

Property of Delaware PT Clinic www.udptclinic.com May be reproduced, as is, for clinical, educational, and research purposes. This Clinical Guideline may need to be modified to meet the needs of a specific patient. The model should not replace clinical judgment.




<p>Weeks 6-10 Dates: _____ to _____</p> <p>Total Visits: 11-24</p> <ul style="list-style-type: none"> D/C use of sling 1-3 visits/wk May begin increasing resistance of flexion week 7 May initiate behind the back IR (towel stretch) week 7 May initiate resistance with ABD week 9 May initiate resistance with ER week 9 	<ul style="list-style-type: none"> <u>Progress ROM to milestones²</u> <ul style="list-style-type: none"> PROM → AAROM → AROM <u>Begin pain-free sub-maximal isometrics at 8 wks³</u> for flexion, extension, ABD, IR, ER <u>Progress to ABD in gravity minimized positions progressing to gravity resisted³</u> <u>Progress to PRE's for all other shoulder motions (flexion, extension, IR)^{3,6}</u> <ul style="list-style-type: none"> Can begin PRE if pain-free with isometrics <u>Progress rhythmic stabilizations^{2,3,6,7}</u> <u>Progress scapular strengthening exercises^{2,3,4,6}</u> <ul style="list-style-type: none"> Progress serratus push-up plus to more horizontal surfaces Bilateral ER/scap retraction in 0° ABD T-band T-Band Rows Prone mid-trap (MT) and low-trap (LT) exercises with scap retraction to plane of the body <u>Modify HEP accordingly</u> 	<ul style="list-style-type: none"> Full A/PROM for all motions except IR in 0° ABD
<p>Weeks 11-12 Dates: _____ to _____</p> <p>Total Visits: 13-30</p> <ul style="list-style-type: none"> 1-3 visits/wk 	<ul style="list-style-type: none"> <u>PRE's for ALL shoulder motions^{2,3,6}</u> <u>Dynamic stabilization exercises^{2,3,6,7}</u> <ul style="list-style-type: none"> Progress rhythmic stabilizations to more challenging and functional positions Shoulder PNF Inertial machine IR/ER beginning in less ABD/ER and progressing to more ABD/ER <u>Continue to progress scapular stabilization/strengthening exercises^{2,3,4,5,6}</u> <ul style="list-style-type: none"> Progress current exercises by increasing resistance/reps/sets Closed chain exercises Quadruped or tripod rhythmic stabilizations Prone LT, MT, HOR ABD Prone rows Standing D2 PNF with T-band <u>PRN: NMES using guidelines at end of protocol⁸</u> <ul style="list-style-type: none"> Infraspinatus starting 8-12 weeks Supraspinatus starting 12-16 weeks 	<ul style="list-style-type: none"> Maintain full A/PROM Independent with HEP Strength improving



<p>Weeks 13-18 Dates: _____ to _____</p> <p>Total Visits: 19-36</p> <ul style="list-style-type: none"> • 1 visit/wk 	<ul style="list-style-type: none"> • <u>Progress strengthening program</u>^{2,3} • <u>Progress dynamic stabilization exercises</u>^{2,3,6,7} <ul style="list-style-type: none"> ○ Progress rhythmic stabilizations to more challenging and functional positions ○ Continue shoulder PNF ○ Continue inertial machine progression • <u>Progress scap strengthening/stabilization exercises</u>^{2,3,4,6} • <u>PRN, NMES to supraspinatus and infraspinatus using guidelines at end of protocol</u>⁸ 	<ul style="list-style-type: none"> • MMT 5/5 all shoulder motions • Full shoulder ROM equal to the uninvolved side
<p>Weeks 19-28 Dates: _____ to _____</p> <ul style="list-style-type: none"> • Physical therapy is as needed for sport/work specific activities 	<ul style="list-style-type: none"> • <u>Continue strengthening and dynamic stabilization exercises as HEP and/or in PT prn</u>^{2,3,6,7} • <u>Begin sport specific interval training program and/or throwing progression</u>^{2,6,7} • <u>Progression of sport/work specific rehabilitation following soreness rules</u>⁹ 	<ul style="list-style-type: none"> • Return to sport/work

Neuromuscular Electrical Stimulation (NMES) Guidelines

- Patient Positioning: seated in a chair with arm in about 30° of elevation in scapular plane and neutral IR/ER using a mobilization belt to prevent movement.
- Electrode Placement
 - Supraspinatus: both pads placed superior to spine of scapula. One pad placed at the medial border of the scapula and one pad placed at lateral border of scapula. Avoid the upper trapezius as much as possible.
 - Infraspinatus: both pads placed inferior to the spine of the scapula. One pad placed at the medial border of the scapula and one pad placed at the lateral border of the scapula.
- Parameters:
 - **EMPI 300PV unit**: Pulse width= 400 microseconds, frequency= 75 pulse per second, on time= 12 seconds, off time= 50 seconds, ramp time= 2 seconds. Intensity to tolerance, goal of visible tetanic contraction.
 - **Versastim**: Pulse width=2500Hz, frequency=75 bursts per second, on time=12 seconds, off time=50 seconds, ramp time=2 seconds. Intensity to tolerance, goal of visible tetanic contraction.





References

1. Morse K, Davis D, Afra R, Kaye EK, Schepesis A, and Voloshin, I. Arthroscopic versus mini-open rotator cuff repair: A comprehensive review and meta-analysis. *Am J Sports Med.* 2008; 36:1824-1828
2. Brotzman SB, and Wilk K. Clinical Orthopaedic Rehabilitation. St. Louis: Mosby, 2003. 125-250.
3. Wilmarth, MA, editor, Davies GJ, Wilk K, Ellenbecker T, Tyler T, Reinhold M, Heiderscheit B, Clark M, Manske R, Matheson J, Kraushaar D, and Mullaley M. *Current Concepts of Orthopaedic Physical Therapy: The Shoulder- Physical Therapy Management utilizing current evidence*, 2nd ed. 2006. APTA independent study course 16.2.4.
4. Moseley JB, Jobe F, Pink M, Perry J, and Tibone J. EMG Analysis of the scapular muscles during a shoulder rehabilitation program. *Am J Sports Med.* 1992;20:128-134.
5. Davies GJ, Dickhoff-Hoffman S. Neuromuscular testing and rehabilitation of the shoulder complex. *JOSPT.* 1993; 18:449-458
6. Kisner C, and Colby LA. Therapeutic Exercise: Foundations and Techniques, 4th ed. Philadelphia: F.A. Davis Company, 2002.
7. Wilk K, Meister K, and Andrews J. Current Concepts in the Rehabilitation of the Overhead Throwing Athlete. *Am J Sports Med.* 2002; 30: 136-151.
8. Reinold MM, Macrina LC, Wilk KE, Dugas JR, Cain EL, Andrews JR. The effect of neuromuscular electrical stimulation of the infraspinatus on shoulder external rotation force production after rotator cuff repair surgery. *Am J Sports Med.* 2008; 36(12): 2317-21
9. Fees M, Decker T, Snyder-Mackler L, Axe MJ. Upper extremity weight training modifications for the injured athlete. A clinical perspective. *Am J Sports Med.* 1998; 26(5): 732-42.

Updated: May 20, 2009