

## PHY 341 HW Ch.2a

Do problems 2.4, 2.5, 2.9; plus the following:

**q2-1** Given the definitions of the lowering and raising operators,

$$a_{\pm} = \frac{\mp ip + m\omega x}{\sqrt{2\hbar m\omega}},$$

show that

$$a_+ a_- = \frac{\hat{H}}{\hbar\omega} - \frac{1}{2}.$$

**q2-2**

Let  $\psi_1(x)$  and  $\psi_2(x)$  be the normalized stationary states with energies  $E_1$  and  $E_2$ , respectively. The wave function  $\Psi(x, 0)$  at  $t = 0$  is given by

$$\Psi(x, 0) = c_1\psi_1(x) + c_2\psi_2(x).$$

- (a) If  $c_1 = \frac{1}{2}$ , find  $c_2$  that makes  $\Psi(x, 0)$  normalized, assuming  $c_2$  is real and positive.
- (b) Find the expectation value of energy. Does it depend on time for  $t > 0$ ?
- (c) Calculate the expectation value of position  $\langle x \rangle$ . Express your answer in terms of  $\bar{x}_1$ ,  $\bar{x}_2$ , and  $\bar{x}_{12}$ , defined as (no need to evaluate)

$$\bar{x}_1 = \int \psi_1^* x \psi_1 dx, \quad \bar{x}_2 = \int \psi_2^* x \psi_2 dx, \quad \bar{x}_{12} = \int \psi_1^* x \psi_2 dx.$$

Does  $\langle x \rangle$  depend on time for  $t > 0$ ?

**q2-3**

Numerical exploration of Problem **q2-2**. Now assume the states are the first two states of a particle in a box. Use the same coefficients  $c_1$  and  $c_2$ , write down of wave function  $\Psi(x, t)$ . Then follow the sample Jupyter code at <http://jwang.sites.umassd.edu/p341/> for superposition of states, plot the probability density  $|\Psi(x, t)|^2$  at different times. Use atomic units (a.u.), where the constants  $m = \hbar = a = 1$ . Argue, based on the graph, that  $\Psi(x, t)$  is not a stationary state, and the expectation value  $\langle x \rangle$  depends on time.

For bonus, plot the real and imaginary parts of  $\Psi(x, t)$ . In Python this can be done as `Psi.real` and `Psi.imag`. Comment on your observations.

Extra credit for the fearless: Numerically check that the normalization  $N = \int_0^a |\Psi(x, t)|^2 dx = 1$  holds for arbitrary  $t$ . Follow the sample code for normalization at the course link. Plot both  $N$  and  $N - 1$  vs time on separate graphs.

You can submit the results and graphs, or email me the code if it contains animation.

**q2-4**

Let  $f = f(x)$  be a differentiable function. By acting each commutator on a test wave function  $\psi$ , show that

- (a)  $[f, x] = 0$ ;
- (b)  $[f, p] = i\hbar f'$ ;
- (c)  $[x, fp] = i\hbar f$ .

<http://jwang.sites.umassd.edu/p341/>