

Newsroom ^(/)

Alzheimer's breaks brain networks' coordination

September 17, 2012

By Michael C. Purdy

Scientists at Washington University School of Medicine (<http://medicine.wustl.edu/>) in St. Louis have taken one of the first detailed looks into how Alzheimer's disease disrupts coordination among several of the brain's networks. The results, reported in *The Journal of Neuroscience*, include some of the earliest assessments of Alzheimer's effects on networks that are active when the brain is at rest.

"Until now, most research into Alzheimer's effects on brain networks has either focused on the networks that become active during a mental task, or the default mode network, the primary network that activates when a person is daydreaming or letting the mind wander," says senior author Beau Ances, MD, assistant professor of neurology. "There are, however, a number of additional networks besides the default mode network that become active when the brain is idling and could tell us important things about Alzheimer's effects."

Ances and his colleagues analyzed brain scans of 559 subjects. Some of these subjects were cognitively normal, while others were in the early stages of very mild to mild Alzheimer's disease. Scientists found that all of the networks they studied eventually became impaired during the initial stages of Alzheimer's.



Ances

"Communications within and between networks are disrupted, but it doesn't happen all at once," Ances says. "There's even one network that has a momentary surge of improved connections before it starts dropping again. That's the salience network, which helps you determine what in your environment you need to pay attention to."

Other networks studied by the researchers included:

- the dorsal attention network, which directs attention toward things in the environment that are salient;
- the control network, believed to be active in consciousness and decision-making; and
- the sensory-motor network, which integrates the brain's control of body movements with sensory feedback (e.g., did the finger that just moved strike the right piano key?).

Scientists also examined Alzheimer's effects on a brain networking property known as anti-correlations. Researchers identify networks by determining which brain areas frequently become active at the same time, but anti-correlated networks are noteworthy for the way their activities fluctuate: When one network is active, the other network is quiet. This ability to switch back and forth between networks is significantly diminished in participants with mild to moderate Alzheimer's disease.

The default mode network, previously identified as one of the first networks to be impaired by Alzheimer's, is a partner in two of the three pairs of anti-correlated networks scientist studied.

"While we can't prove this yet, one hypothesis is that as things go wrong in the processing of information in the default mode network, that mishandled data is passed on to other networks, where it creates additional problems," Ances says.

It's not practical to use these network breakdowns to clinically diagnose Alzheimer's disease, Ances notes, but they may help track the development of the disease and aid efforts to better understand its spread through the brain.

Ances plans to look at other markers for Alzheimer's disease in the same subjects, such as levels in the cerebrospinal fluid of amyloid beta, a major component of Alzheimer's plaques.

Brier MR, Thomas JB, Snyder AZ, Benziger TL, Zhang D, Raichle ME, Holtzman DM, Morris JC, Ances BM. Loss of intranetwork and internetwork resting state functional connections with Alzheimer's progression. *The Journal of Neuroscience*, June 27, 2012. DOI:10.1523/JNEUROSCI.5698-11.2012

This research was supported by funding from the Medical Scientist Training Program Grant to Washington University School

of Medicine (5T32GM007200-37); a Knight Alzheimer's Disease Research Center Pilot Grant (3255 ADRC 26); the National Institute of Mental Health (K23MH081786); the National Institute of Nursing Research (RO1NR012907 and RO1NR012657); the Dana Foundation; the Alene and Meyer Kopolov Fund for Geriatric Psychiatry and Neurology; the National Institute of Aging (PO1AG026276, PO1AG026276, PO1AG03991, and P50 AG05681); the National Institute of Neurological Disorders and Stroke (NS06833); the Washington University Center for Translational Neuroscience (P30NS057105); and the American Roentgen Ray Society Foundation.

Washington University School of Medicine (<http://medicine.wustl.edu/>)'s 2,100 employed and volunteer faculty physicians also are the medical staff of Barnes-Jewish (<http://www.barnesjewish.org/>) and St. Louis Children's (<http://www.stlouischildrens.org/>) hospitals. The School of Medicine is one of the leading medical research, teaching and patient care institutions in the nation, currently ranked sixth in the nation by *U.S. News & World Report*. Through its affiliations with Barnes-Jewish and St. Louis Children's hospitals, the School of Medicine is linked to BJC HealthCare (<http://www.bjc.org/>).