

The Role of Criterion A2 in the DSM-IV Diagnosis of Posttraumatic Stress Disorder

Elie George Karam, Gavin Andrews, Evelyn Bromet, Maria Petukhova, Ayelet Meron Ruscio, Mariana Salamoun, Nancy Sampson, Dan J. Stein, Jordi Alonso, Laura Helena Andrade, Matthias Angermeyer, Koen Demyttenaere, Giovanni de Girolamo, Ron de Graaf, Silvia Florescu, Oye Gureje, Debra Kaminer, Roman Kotov, Sing Lee, Jean-Pierre Lépine, Maria Elena Medina-Mora, Mark A. Oakley Browne, José Posada-Villa, Rajesh Sagar, Arieh Y. Shalev, Tadashi Takeshima, Toma Tomov, and Ronald C. Kessler

Background: Controversy exists about the utility of DSM-IV posttraumatic stress disorder (PTSD) criterion A2 (A2): that exposure to a potentially traumatic experience (PTE; PTSD criterion A1) is accompanied by intense fear, helplessness, or horror.

Methods: Lifetime DSM-IV PTSD was assessed with the Composite International Diagnostic Interview in community surveys of 52,826 respondents across 21 countries in the World Mental Health Surveys.

Results: Of 28,490 representative PTEs reported by respondents, 37.6% met criterion A2, a proportion higher than the proportions meeting other criteria (B–F; 5.4%–9.6%). Conditional prevalence of meeting all other criteria for a diagnosis of PTSD given a PTE was significantly higher in the presence (9.7%) than absence (.1%) of A2. However, as only 1.4% of respondents who met all other criteria failed A2, the estimated prevalence of PTSD increased only slightly (from 3.64% to 3.69%) when A2 was not required for diagnosis. Posttraumatic stress disorder with or without criterion A2 did not differ in persistence or predicted consequences (subsequent suicidal ideation or secondary disorders) depending on presence-absence of A2. Furthermore, as A2 was by far the most commonly reported symptom of PTSD, initial assessment of A2 would be much less efficient than screening other criteria in quickly ruling out a large proportion of noncases.

Conclusions: Removal of A2 from the DSM-IV criterion set would reduce the complexity of diagnosing PTSD, while not substantially increasing the number of people who qualify for diagnosis. Criterion A2 should consequently be reconceptualized as a risk factor for PTSD rather than as a diagnostic requirement.

Key Words: Composite International Diagnostic Interview (CIDI), criterion A2, diagnosis, DSM-IV, posttraumatic stress disorder (PTSD), potentially traumatic experience (PTE), World Health Organization World Mental Health (WMH) Surveys

Controversy exists over what should qualify as a trauma for posttraumatic stress disorder (PTSD). Some experts favor a broader and others a narrower definition (1,2). The DSM-IV (3) tries to find a middle ground by including both an objective

(criterion A1 [A1]) and a subjective (criterion A2 [A2]) component. Criterion A1 is broader than favored by some, as it includes any event a person “experienced, witnessed, or (was) confronted with” that involves “actual or threatened death or serious injury or a threat to the physical integrity of oneself or others.” The inclusion of indirect exposure and ambiguity of the term “threat to physical integrity” are particular points of controversy (4,5). Criterion A2, new to DSM-IV, was designed to restrict qualifying events to those where the immediate response involved “intense fear, helplessness,

From the Department of Psychiatry and Clinical Psychology (EGK, MS), Saint George Hospital, University Medical Center; Department of Psychiatry and Clinical Psychology, Faculty of Medicine, Balamand University Medical School; and the Institute for Development, Research, Advocacy and Applied Care, Beirut, Lebanon; Department of Psychiatry (GA), University of New South Wales, Sydney, Australia; State University of New York (EB), Stony Brook, New York; Department of Health Care Policy (MP, NS, RCK), Harvard Medical School, Boston, Massachusetts; Department of Psychology (AMR), University of Pennsylvania, Philadelphia, Pennsylvania; Department of Psychiatry (DJS), Groote Schuur Hospital, Cape Town, South Africa; Health Services Research Unit (JA), Institut Municipal d'Investigació Mèdica, Hospital Del Mar, Centro de Investigación Biomédica en Red en Epidemiología y Salud Pública, Barcelona, Spain; Section of Psychiatric Epidemiology-LIM 23 Department and Institute of Psychiatry (LHA), School of Medicine, University of São Paulo, São Paulo, Brazil; Center for Public Mental Health (MA), Gössing am Wagram, Austria; Department of Psychiatry (KD), University Hospital Gasthuisberg, Leuven, Belgium; Istituto de Ricovero e Cura a Carattere Scientifico Centro S. Giovanni di Dio Fatebenefratelli (GdG), Brescia, Italy; Monitoring & Epidemiology Department, Psychiatric Epidemiology Unit, Netherlands Institute of Mental Health and Addiction (RdG), Utrecht, The Netherlands; Public Health Research and Evidence Based Medicine Department (SF), Department of Psychiatry, National School of Public Health and Health Services Management, Bucharest, Romania; University College Hospital (OG), Prince of Wales, Ibadan, Nigeria; MRC Research Unit on Anxiety Disorders, University of Stellenbosch (DK), Cape Town, South Africa; Department of Psychiatry (RK), Stony Brook University, Stony Brook, New York; Department of Psychiatry (SL), The Chinese University of Hong Kong, Hong Kong, China; Institut national de la santé et de la recherche médicale, U705, Centre national de la recherche scientifique Unité Mixte de Recherche 7157 (JPL), University Paris Diderot, Hôpital Lariboisière Fernand Widal, Assistance Publique Hôpitaux de Paris, Paris France; National Institute of Psychiatry Ramon de la Fuente (MEMM), Mexico City, Mexico; the Discipline of Psychiatry (MAOB), The University of Tasmania, Tasmania, Australia; Colegio Mayor de Cundinamarca University (JP-V), Bogota, Columbia; Department of Psychiatry (RS), All India Institute of Medical Sciences, New Delhi, India; Department of Psychiatry (AYS), Hadassah University Hospital, Jerusalem, Israel; National Institute of Mental Health (TTa), National Center of Neurology and Psychiatry, Tokyo, Japan; and Department of Psychiatry (TTo), New Bulgarian University, Sofia, Bulgaria.

Address correspondence to Ronald C. Kessler, Ph.D., Department of Health Care Policy, Harvard Medical School, 180 Longwood Avenue, Boston, MA 02115; E-mail: kessler@hcp.med.harvard.edu.

Received Dec 22, 2009; revised Apr 30, 2010; accepted Apr 30, 2010.

Table 1. World Mental Health Sample Characteristics by World Bank Income Categories

Country by Income Category ^a	Survey	Sample Characteristics ^b	Field Dates	Age Range	Sample Size		Response Rate ^c
					Part I	Part II	
I. High-Income Countries							
Belgium	ESEMeD	Stratified multistage clustered probability sample of individuals residing in households from the national register of Belgium residents. NR	2001–2002	18+	2,419	1,043	50.6
France	ESEMeD	Stratified multistage clustered sample of working telephone numbers merged with a reverse directory (for listed numbers). Initial recruitment was by telephone, with supplemental in-person recruitment in households with listed numbers. NR	2001–2002	18+	2,894	1,436	45.9
Germany	ESEMeD	Stratified multistage clustered probability sample of individuals from community resident registries. NR	2002–2003	18+	3,555	1,323	57.8
Israel	NHS	Stratified multistage clustered area probability sample of individuals from a national resident register. NR	2002–2004	21+	4,859	4,859	72.6
Italy	ESEMeD	Stratified multistage clustered probability sample of individuals from municipality resident registries. NR	2001–2002	18+	4,712	1,779	71.3
Japan	WMHJ 2002–2006	Unclustered two-stage probability sample of individuals residing in households in 11 metropolitan areas.	2002–2006	20+	4,129	1,682	55.1
Netherlands	ESEMeD	Stratified multistage clustered probability sample of individuals residing in households that are listed in municipal postal registries. NR	2002–2003	18+	2,372	1,094	56.4
New Zealand ^d	NZMHS	Stratified multistage clustered area probability sample of household residents. NR	2003–2004	18+	12,790	7,312	73.3
Spain	ESEMeD	Stratified multistage clustered area probability sample of household residents. NR	2001–2002	18+	5,473	2,121	78.6
United States Total	NCS-R	Stratified multistage clustered area probability sample of household residents. NR	2002–2003	18+	9,282	5,692	70.9
					52,485	28,341	
II. Upper-Middle-Income Countries							
Brazil	São Paulo Megacity	Stratified multistage clustered area probability sample of household residents in the São Paulo metropolitan area.	2005–2007	18+	5,037	2,942	81.3
Bulgaria	NSHS	Stratified multistage clustered area probability sample of household residents. NR	2003–2007	18+	5,318	2,233	72.0
Lebanon	LEBANON	Stratified multistage clustered area probability sample of household residents. NR	2002–2003	18+	2,857	1,031	70.0
Mexico	M-NCS	Stratified multistage clustered area probability sample of household residents in all urban areas of the country (approximately 75% of the total national population).	2001–2002	18–65	5,782	2,362	76.6
Romania	RMHS	Stratified multistage clustered area probability sample of household residents. NR	2005–2006	18+	2,357	2,357	70.9
South Africa	SASH	Stratified multistage clustered area probability sample of household residents. NR	2003–2004	18+	4,315	4,315	87.1
Total					25,666	15,240	
III. Low-Income/Lower-Middle-Income Countries							
Colombia	NSMH	Stratified multistage clustered area probability sample of household residents in all urban areas of the country (approximately 73% of the total national population).	2003	18–65	4,426	2,381	87.7
India	WMHI	Stratified multistage clustered area probability sample of household residents in Pondicherry region. NR	2003–2005	18+	2,992	1,373	98.8
Nigeria	NSMHW	Stratified multistage clustered area probability sample of households in 21 of the 36 states in the country, representing 57% of the national population. The surveys were conducted in Yoruba, Igbo, Hausa, and Efik languages.	2002–2003	18+	6,752	2,143	79.3
People's Republic of China	B-WMH S-WMH	Stratified multistage clustered area probability sample of household residents in the Beijing and Shanghai metropolitan areas.	2002–2003	18+	5,201	1,628	74.7
Ukraine	CMDPSD	Stratified multistage clustered area probability sample of household residents. NR	2002	18+	4,725	1,720	78.3

Table 1. (continued)

Country by Income Category ^a	Survey	Sample Characteristics ^b		Field Dates	Age Range	Sample Size		Response Rate ^c
		Part I	Part II			Part I	Part II	
Total		24,096	9,245			24,096	9,245	
IV. Total		102,247	52,826			102,247	52,826	

B-WMH, The Beijing World Mental Health Survey; CMDPSD, Comorbid Mental Disorders During Periods of Social Disruption; ESEMEd, The European Study of the Epidemiology of Mental Disorders; LEBANON, Lebanese Evaluation of the Burden of Ailments and Needs Of the Nation; M-NCS, The Mexico National Comorbidity Survey; NCS-R, The US National Comorbidity Survey Replication; NHS, Israel National Health Survey; NR, nationally representative; NSHS, Bulgaria National Survey of Health and Stress; NSMH, The Colombian National Study of Mental Health; NSMHW, The Nigerian Survey of Mental Health and Wellbeing; NZMHS, New Zealand Mental Health Survey; RMHS, Romania Mental Health Survey; SASH, South Africa Health Survey; S-WMH, The Shanghai World Mental Health Survey; WMHI, World Mental Health India; WMHJ2002–2006, World Mental Health Japan Survey.

^aThe World Bank (2008). Data and Statistics. Accessed May 12, 2009 at: <http://go.worldbank.org/D75N0B8YUO>.

^bMost World Mental Health surveys are based on stratified multistage clustered area probability household samples in which samples of areas equivalent to counties or municipalities in the United States were selected in the first stage followed by one or more subsequent stages of geographic sampling (e.g., towns within counties, blocks within towns, households within blocks) to arrive at a sample of households, in each of which a listing of household members was created and one or two people were selected from this listing to be interviewed. No substitution was allowed when the originally sampled household resident could not be interviewed. These household samples were selected from Census area data in all countries other than France (where telephone directories were used to select households) and the Netherlands (where postal registries were used to select households). Several World Mental Health surveys (Belgium, Germany, Italy) used municipal registries to select respondents without listing households. The Japanese sample is the only totally unclustered sample, with households randomly selected in each of the 11 sample areas and one random respondent selected in each sample household. Sixteen of the 22 surveys are based on nationally representative household samples, while two others are based on nationally representative household samples in urbanized areas (Colombia, Mexico).

^cThe response rate is calculated as the ratio of the number of households in which an interview was completed to the number of households originally sampled, excluding from the denominator households known not to be eligible either because of being vacant at the time of initial contact or because the residents were unable to speak the designated languages of the survey. The weighted average response rate is 71.6%.

^dNew Zealand interviewed respondents 16+ but for the purposes of cross-national comparisons we limit the sample to those 18+.

or horror” to prevent the overdiagnosis that might otherwise occur because of the broad scope of A1.

The few studies that focus specifically on A2 show a substantial proportion of people exposed to an A1 experience—a potentially traumatic experience (PTE)—meets A2 (4,6). This means the A2 requirement reduces estimated prevalence only modestly (1,7). However, failure to meet A2 strongly predicts failure to meet other criteria (6), making assessment of A2 useful in two ways. First, as A2 is the only PTSD criterion that can be assessed near the time of PTE exposure, early assessment of A2 can help predict subsequent PTSD, although recent prospective research shows that a meaningful proportion of patients who subsequently meet all other criteria for PTSD fail to meet A2 at the time of trauma exposure (8). Second, even when the assessment takes place months or years after PTE exposure, initial retrospective assessment of A2 can provide a quick rule-out (5). At the same time, the fact that PTSD diagnoses would increase only modestly if A2 was removed as a requirement argues against retaining A2 based on the interest in simplifying diagnoses by eliminating redundant criteria (9,10).

The empiric foundation on which these competing views are based is limited to a small number of United States and Australian studies. Only scant data exist, furthermore, on the relationship between A2 and other clinical correlates (4), making it unclear whether the A2 requirement excludes clinically significant cases. Given increasing adoption of DSM-IV criteria worldwide, it is important to evaluate these issues using cross-national data (11). We do this here with data from the World Health Organization (WHO) World Mental Health (WMH) Survey Initiative (12).

Methods and Materials

Samples

Data come from WMH surveys in 21 countries, 5 low-income or lower-middle-income (Colombia, India, Nigeria, People's Republic of China [Beijing and Shanghai], and Ukraine), 6 upper-middle-income (Brazil, Bulgaria, Lebanon, Mexico, Romania, and South Africa), and 10 high-income (Belgium, France, Germany, Israel, Italy, Japan, Netherlands, New Zealand, Spain, and United States) countries (Table 1). A total of 102,247 respondents were interviewed in these surveys. Part I of the interviews, which assessed core disorders, was completed by all respondents. Part II assessed additional disorders, including PTSD, and correlates and was completed by 100% of respondents who met criteria for any part I disorder and a probability subsample of other part I respondents. Part II sample sizes range between 1031 (Lebanon) and 7312 (New Zealand) and total 52,826 respondents across countries. This is the sample used here. All surveys used multistage clustered area probability household samples representative of specific regions (Colombia, Japan, Mexico, Nigeria, People's Republic of China) or entire nations (the remaining countries). The weighted (by sample size) average response rate was 71.6%. The part I sample was weighted to adjust for differential probabilities of selection and residual discrepancies between sample and census on sociodemographic and geographic variables. The part II sample was additionally weighted to adjust for undersampling of part I respondents without part I disorders. World Mental Health sampling and weighting are discussed in more detail elsewhere (13).

Procedures

Respondents were interviewed face-to-face by trained lay interviewers who obtained informed consent before initiating

interviews. Recruitment-consent procedures were approved by human subjects committees in each country. Interviewer training and quality control procedures were cross-nationally standardized (14). The instrument was the WHO Composite International Diagnostic Interview (CIDI) (15), a fully structured interview. Translation and back translation followed standard WHO procedures (16).

The Assessment of DSM-IV PTSD

The CIDI PTSD section began with questions about lifetime occurrence of 27 PTEs (Table 2). An additional open-ended question then asked about any other traumatic event. Positive responses were recorded verbatim and reviewed by a clinical rater to confirm PTSD criterion A1. A second open-ended question then attempted to obtain information about qualifying events that respondents did not report because of embarrassment. Wording was as follows: “[s]ometimes people have experiences they don’t want to talk about in interviews. I won’t ask you to describe anything like this, but without telling me what it was, did you ever have a traumatic event that you did not report to me because you did not want to talk about it?”

Table 2. The Potentially Traumatic Experiences Assessed in the World Mental Health Surveys

-
- I. Experiences Involving Interpersonal Violence
 - 1. Combat experience (military or nonmilitary) in a war or sectarian violence (e.g., political, religious, or ethnic conflicts)
 - 2. Relief worker or peacemaker in war zone or region of sectarian violence
 - 3. Civilian in a war zone
 - 4. Civilian in a region of sectarian violence
 - 5. Displaced refugee from a war zone or a region of sectarian violence
 - 6. Kidnapped or held captive
 - 7. Beaten up as a child by a caregiver
 - 8. Beaten up by a spouse or romantic partner
 - 9. Beaten up by someone else
 - 10. Mugged or threatened with a weapon
 - 11. Raped
 - 12. Sexually assaulted or molested
 - 13. Stalked
 - II. Other Threats to the Physical Integrity of the Respondent
 - 14. Natural disaster (e.g., flood, hurricane, earthquake)
 - 15. Life-threatening automobile accident
 - 16. Other life-threatening accident
 - 17. Toxic chemical exposure
 - 18. Other exposure to a made disaster (e.g., fire, explosion at a place of work)
 - 19. Other life-threatening illness or injury
 - III. Threats to the Physical Integrity of Others
 - 20. Death of a loved one
 - 21. Life-threatening illness of a loved one
 - 22. Any other trauma experienced by a loved one
 - 23. Witnessed repeated physical fights at home as a child
 - 24. Witnessed any other injury or death
 - 25. Accidentally caused injury or death to someone
 - 26. Purposefully injured, tortured, or killed someone
 - 27. Witnessed atrocities or carnage
 - IV. Open-Ended Questions About Other Potentially Traumatic Experiences
 - 28. Any other objectively qualifying experiences (respondents are asked to describe these experiences)
 - 29. Private experiences (respondents are explicitly told in advance that they will not be asked to describe these experiences)
-

As we knew from previous research (6,17) that some respondents report too many PTEs to assess PTSD for each one, we assessed one randomly selected PTE plus the respondent’s self-nominated worst lifetime PTE. The random event is the focus of the current report, as appropriate weighting yields a representative sample of all PTEs. We sought such a sample because the conventional approach of focusing only on the worst PTE gives a biased perspective on event exposure and impact (18). However, we also report results for PTEs that received treatment to ensure that results apply to clinically relevant PTEs.

The random event was selected by numbering each PTE and either having the computer select one random number from those endorsed (in countries that used computer-assisted interviewing) or assigning a random start value unique to each respondent and selecting the first endorsed event after that value (in countries that used paper-and-pencil interviewing). The latter method does not guarantee equal probability of selection of PTEs, requiring an additional weight to adjust for differences in probability of selection as a function of placement in the list. Once the random event type was selected, a random occurrence of that event was selected when there were multiple occurrences. We then weighted the data by the number of unique occurrences of all qualifying PTEs to produce a weighted dataset in which each PTE is represented in the proportion it occurred in the population. As an indication of PTE distribution, the mean number of PTE occurrences among respondents with more than one was 5.8, with a range of 2 to 160 and interquartile range of 3 to 6.

Criterion A2 was considered met if the respondent endorsed any of three questions about whether, at the time of the random PTE, he or she felt 1) terrified or very frightened, 2) helpless, or 3) shocked or horrified. The remaining criteria were then assessed as whether or not A2 was endorsed with questions about re-experiencing (criterion B), avoidance/numbing (criterion C), arousal (criterion D), duration (criterion E), and clinically significant distress impairment (criterion F). A retrospective question asked respondents how many months or years symptoms continued. All responses were converted into months (e.g., 5 years = 60 months) for analysis of persistence. Blinded clinical reappraisal interviews with the Structured Clinical Interview for DSM-IV (SCID) (19) carried out in the United States WMH survey (20) found an area under the receiver operating characteristic curve of .69 between PTSD diagnoses based on the CIDI and SCID.

The Assessment of Correlates of PTSD

Other lifetime DSM-IV/CIDI disorders considered were anxiety disorders (panic disorder, phobias, generalized anxiety disorder), mood disorders (major depressive disorder, dysthymic disorder, bipolar I–II disorder), and substance disorders (alcohol and drug abuse or dependence with abuse). Organic exclusion rules and diagnostic hierarchy rules were used in making these diagnoses (15). As described elsewhere (21), blinded SCID clinical reappraisal interviews (19) in the United States WMH sample documented generally good CIDI-SCID concordance (area under the receiver operating characteristic curve) for diagnoses of anxiety (.73), mood (.93), substance (.86), and any (.76) disorder. Lifetime suicidal ideation was assessed with a single question that asked respondents if they ever seriously thought about killing themselves and, if so, their age when this first occurred.

Analysis Methods

Cross-tabulations examined frequency of individual PTEs, conditional prevalence of DSM-IV PTSD with and without A2,

and co-occurrence of criteria B through F. Logistic regression (22) examined sociodemographic predictors of PTSD with or without criterion A2 and interactions of predictors with presence-absence of A2. Discrete-time survival analysis (23) with person year the unit of analysis was used to examine association between PTSD with and without A2 and persistence, as well as subsequent first onset of other DSM-IV/CIDI disorders and suicidal ideation. All parameters were estimated using the Taylor series method (24) in the SUDAAN software system (Research Triangle Institute, Research Triangle Park, North Carolina) (25). Significance was evaluated using .05 level two-sided Wald χ^2 tests.

Results

Prevalence of DSM-IV PTSD Criteria A1 and A2

Lifetime exposure to PTEs was reported by 67.9% of respondents (72.8% in high-income, 61.7% in upper-middle-income, and 63.1% in low-income/lower-middle-income countries). Disaggregated comparisons (detailed results available on request) found the higher exposure in high-income countries due largely to automobile accidents. Weighted mean number of PTEs per respondent with any was 4.5 (4.7 in high-income, 4.1 in upper-middle-income, and 4.6 in low-income/lower-middle-income countries) for 305.6 lifetime PTEs for every 100 respondents (i.e., $67.9\% \times 4.5$) and 137,778 PTEs in the full sample.

Other Criteria for DSM-IV PTSD

Approximately one third (37.6%) of weighted PTEs were reported as meeting A2 (Table 3). This proportion is substantially lower in high-income than upper-middle-income or low-income/lower-middle-income countries. Conditional prevalence of meeting all other DSM-IV requirements for PTSD is dramatically higher in the presence (9.7%) than absence (.1%) of A2. Conditional prevalence of PTSD given A2 is positively related to country income level. Conditional prevalence of all other requirements for PTSD in the absence of A2 is very low (.0%–.1%) regardless of country income level.

It is unclear whether these results, which involve retrospective reporting of A2 in some cases many years after the PTE, accurately reflect the way the respondent would have characterized their short-term reactions at the time of the PTE. To the

extent that they do, though, the results would mean that an assessment of A2 shortly after PTE occurrence might help predict subsequent PTSD. Indeed, if we compare the proportion of all random PTEs that meet full criteria for PTSD (by multiplying conditional prevalence of PTSD given A2 by the proportion of all PTEs with A2, both reported in Table 3) with the additional proportion that would meet criteria for PTSD if A2 was no longer required for diagnosis (by multiplying conditional prevalence of PTSD without A2 by the proportion of all PTEs without A2, both reported in Table 3), we find that the retrospective results suggest that only 1.4% of respondents who met all other requirements for PTSD would be missed if follow-up assessment was carried out only among people with A2. This estimated proportion is very similar in high-income, upper-middle-income, and low-income/lower-middle-income countries (.6%–2.0%). More detailed examination (available on request) found no country or PTE for which the proportion of respondents meeting all other requirements who would be missed by focusing on A2-positives is significantly more than 2%. That is, the upper end of the 95% confidence interval of these prevalence estimates never exceeded 2%. Furthermore, additional analyses (detailed results available on request) found that the percentage is even smaller (1.1%) among PTEs for which professional treatment was sought, demonstrating that the low percentage holds for the most clinically relevant PTEs.

As noted in the introduction, one other rationale for including A2 in DSM-IV was that it provided a quick rule-out for noncases. As it happens, though, our results suggest this is not true. Exclusion of PTEs that failed A2 would reduce by an estimated 62.4% (i.e., $100\% - \text{the } 37.6\% \text{ of PTEs that met criterion A2}$) the number who would have to be further assessed and would miss only 1.4% who met all other criteria. Although this seems like a very effective screening rule, higher conditional prevalence of A2 than other DSM-IV criteria means the proportion of noncases ruled out by A2 would be considerably lower than those ruled out by assessing any other DSM-IV criterion. For example, assessing criterion F (clinically significant distress-impairment) would exclude 90.4% of noncases compared with 62.4% by assessing A2.

Table 3. Percentage of the Representative Sample of Potentially Traumatic Experiences Assessed in the World Mental Health Sample That Had Criterion A2 and Conditional Prevalence of DSM-IV/Composite International Diagnostic Interview Posttraumatic Stress Disorder in the Presence Versus Absence of Criterion A2

Country Income	Percentage of PTEs That Had Criterion A2			Conditional Prevalence of PTSD in the Presence of Criterion A2			Conditional Prevalence of All Other Requirements for a Diagnosis of PTSD in the Absence of Criterion A2		
	%	(SD)	(n) ^a	%	(SD)	(n) ^a	%	(SD)	(n) ^a
Levels									
All	37.6	(.9)	(28,490)	9.7	(.7)	(11,121)	.1	(.0)	(17,369)
High	28.6	(1.1)	(17,815)	15.0	(1.2)	(6,341)	.1	(.0)	(11,474)
Upper-middle	52.2	(1.7)	(6,962)	6.3	(.8)	(3,103)	.1	(.1)	(3,859)
Low/lower-middle	49.5	(2.7)	(3,713)	3.8	(1.0)	(1,677)	.0	(.0)	(2,036)

PTE, potentially traumatic experience; PTSD, posttraumatic stress disorder.

^aThe *n*'s reported here are unweighted denominators of the number of potentially traumatic experiences (PTEs) on which the prevalence estimates are based. For example, a weighted 36.7% of the 28,490 PTEs reported (an observed 11,121 PTEs) were classified as meeting the DSM-IV/Composite International Diagnostic Interview A2 criterion, while a weighted 9.7% of these 11,121 PTEs were classified as meeting all other DSM-IV/Composite International Diagnostic Interview criteria for posttraumatic stress disorder. The ratio of the number of respondents who reported at least one PTE to the total sample size ($28,490/52,949 = 53.8\%$) is lower than the weighted 67.9% of respondents who reported a PTE because people with high rates of PTE exposure were somewhat underrepresented in the samples.

The Clinical Significance of PTSD Without A2

The above results show that removal of A2 from the criterion set would have very little effect on the number of people who qualify for a diagnosis of PTSD based on randomly selected PTEs in the WMH sample. The PTSD prevalence based on randomly selected PTEs would increase only from 3.64% to 3.69% (about a 2% increase on the base of 3.64%) if we excluded A2, adding only 21 cases classified as meeting full criteria for PTSD out of the 28,490 cases in the weighted sample of randomly selected PTEs (.07% of all such PTEs). In considering whether such a modest difference justifies inclusion of A2 in the DSM-IV criterion set, it is relevant to know whether the few otherwise qualifying cases are so distinct in terms of low clinical significance that they need to be excluded. We addressed this question by examining whether PTSD differs in course depending on presence-absence of A2. No such difference was found. The odds ratio (OR) (95% confidence interval in parentheses) of presence-absence of A2 predicting differential persistence of PTSD among cases with versus without A2 was insignificant [1.02 (.42–2.47), $\chi^2_1 = .0, p = .97$]. The OR of PTSD without A2 predicting subsequent first onset of other DSM-IV/CIDI disorders was actually higher, although not significantly so ($\chi^2_1 = 1.0, p = .32$), for PTSD without than with A2: 9.9 (2.1–47.1) versus 4.4 (3.2–6.1). The same basic pattern holds in predicting subsequent onset of suicidal ideation with an OR (95% confidence interval) of 3.6

(2.8–4.7) for PTSD with A2 and 4.4 (.8–24.7) without A2 ($\chi^2_1 = .1, p = .83$).

Compositional Differences Between PTSD With and Without A2

We investigated whether the composition of cases of PTSD differs depending on presence-absence of A2 using logistic regression where information about age at time of PTE exposure, sex, history of prior lifetime DSM-IV/CIDI disorders, and history of prior PTE exposure were used to distinguish PTSD with and without A2 (Table 4). Men had significantly higher odds than women of PTSD without than with A2 (5.1). Respondents with a history of other DSM-IV/CIDI disorders had significantly lower odds than others of PTSD without than with A2 (.2).

Discussion

This study found that PTEs are commonly occurring, that a much higher proportion of the randomly selected PTEs meet criterion A2 than any other DSM-IV PTSD criterion, that conditional prevalence of meeting diagnostic threshold based on other criteria is significantly higher in the presence than absence of A2, and that only a small fraction of respondents who meet diagnostic threshold based on other criteria fail to meet A2. The last of these results means the sample-specific prevalence of PTSD increased only very slightly when A2 was not required for

Table 4. Logistic Regression Equations Predicting Whether the Potentially Traumatic Experience Met Criterion A2 in the Sample of Potentially Traumatic Experiences Assessed in the World Mental Health Sample That Satisfy All Other Criteria for a DSM-IV/Composite International Diagnostic Interview Diagnosis of Posttraumatic Stress Disorder

	Country Income Level							
	All		High		Upper-Middle		Low/Lower-Middle	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Sex								
Male	5.1 ^a	(1.6–16.5)	8.4 ^a	(2.1–33.8)	.8	(.1–4.9)	12.1 ^a	(1.1–129.4)
Female	1.0	–	1.0	–	1.0	–	1.0	–
χ^2_1		7.4 ^a		9.0 ^a		.1		4.2 ^a
Age at Trauma Exposure								
0–9	.5	(.1–3.3)	.8	(.1–6.9)	.0	(.0–.3)	– ^c	–
10–19	.7	(.1–5.0)	.9	(.1–9.6)	.4	(.0–6.5)	–	–
20–29	.5	(.1–3.7)	.6	(.1–6.8)	.1	(.0–.9)	–	–
30–39	1.7	(.3–12.0)	1.2	(.1–21.8)	1.0	(.1–13.6)	–	–
40+	1.0	–	1.0	–	1.0	–	–	–
χ^2_4		1.9		.4		8.9		– ^c
History of Prior (to Age of PTE Exposure) DSM-IV/CIDI Disorders								
Yes	.2 ^a	(.1–1.0)	.2	(.0–1.5)	.4	(.0–3.4)	.3	(.1–1.1)
No	1.0	–	1.0	–	1.0	–	1.0	–
χ^2_1		4.0 ^a		2.4		.8		3.1
History of Prior (to Age of Occurrence of the Random PTE) PTE Exposure								
Yes	.8	(.2–2.9)	1.0	(.2–5.8)	.5	(.1–2.4)	1.1	(.1–9.1)
Number	1.0	–	1.0	–	1.0	–	1.0	–
χ^2_1		.2		.0		.9		.0
(n)		(1,340) ^b		(981) ^b		(267) ^b		(92) ^b

CI, confidence interval; CIDI, Composite International Diagnostic Interview; OR, odds ratio; PTE, potentially traumatic experience.

^aSignificant at the .05 level, two-sided test.

^bThe *n*'s reported here are the unweighted numbers of potentially traumatic experiences (PTEs) on which the regression equations are based. These include all PTEs in the representative sample of PTEs that were judged to meet all other DSM-IV/Composite International Diagnostic Interview criteria for a diagnosis of posttraumatic stress disorder in the presence or absence of criterion A2. The outcome is whether or not the PTE met all other posttraumatic stress disorder criteria but not criterion A2.

^cThe number of observed cases of posttraumatic stress disorder without criterion A2 in this subsample was too small to estimate the association with this polytomous variable.

diagnosis. Furthermore, cases meeting all other PTSD criteria do not differ in persistence, subsequent suicidal ideation, or secondary disorders from PTSD cases. We concluded from these results that A2 should be reconceptualized as a risk factor for PTSD rather than as a diagnostic requirement, as the retrospective data indirectly suggest that the occurrence of A2 symptoms shortly after the occurrence of the PTE is significantly predictive of subsequent PTSD, even though the inclusion of A2 as a requirement for diagnosis has little effect on case definition once other diagnostic requirements are met.

Our finding that a substantial minority of PTEs was reported as causing intense fear, helplessness, or horror in countries throughout the world is consistent with previous studies in a narrower range of countries (4,6,7,26). It is unclear, though, why conditional prevalence of A2 is significantly lower in high-income (28.6%) than lower-income (49.5%–52.2%) countries. This could be due to PTEs in lower-income countries being, on average, more extreme (either in terms of characteristics of or objective consequences of the events) than in high-income countries. Another possibility is that people in low-income countries are more likely than those in high-income countries to be shocked or horrified by the occurrence of a traumatic stress. We are not aware of any cross-national research that addresses this second possibility, but it is indirectly inconsistent with research in the United States that documented stronger effects of some traumatic events in leading to shock and horror among middle class than lower class people due to middle class people being more likely to have illusory world views about control and justice that are shattered by exposure to traumatic events (27,28). Why we would find the opposite pattern in cross-national comparisons is perplexing. Another significant between-country difference is that conditional prevalence of PTSD given A2 is significantly higher in high-income (15.0%) than lower-income (3.8%–6.3%) countries. These differences might reflect between-country differences in coping resources that influence emotional reactions to trauma.

Our finding that conditional prevalence of A2 given A1 far exceeded conditional prevalence of any other DSM-IV criterion of PTSD suggests that it would be more efficient to use a criterion other than A2 for screening to quickly rule out a diagnosis of PTSD. This is due to the high prevalence of A2 occurring not only among people with PTSD but also among people without PTSD.

Our finding that only a small proportion of PTEs without A2 were associated with all other DSM-IV requirements for PTSD is consistent with most (1,4,6,7), but not all (8), previous studies. This means that removal of A2 from the criterion set would lead to only a modest increase in the number of people diagnosed with PTSD. Furthermore, the small number of cases of PTSD without A2 in the WMH series had equal PTSD persistence and equally elevated risk of temporally secondary mental disorders and suicidal ideation as cases with A2. This means that the small number of otherwise qualifying cases excluded by A2 should not be excluded. This conclusion is reinforced by evidence from other research that A2 is often absent in patients from high-risk occupations (e.g., military, police, fire fighters) who otherwise meet criteria for PTSD associated with traumatic events that occurred in the context of their occupation (29), as well as among those who were amnesic (8).

An especially intriguing finding was that conditional risk of PTSD without versus with A2 was significantly higher among respondents with than without a history of other DSM-IV/CIDI disorders. This might have been caused by prior psychopathology creating stronger vulnerability for PTSD associated with less

severe than more severe PTEs, where the occurrence of A2 symptoms is a marker of PTE severity. This possibility is consistent with the general diathesis-stress model of anxiety disorders (30). This model suggests that increased vulnerability to stress (as indicated here by history of psychopathology) reduces the amount of stress needed to promote onset of a stress-related anxiety disorder, whereas vulnerability factors become less important at high levels of stress exposure due to resistance resources becoming overwhelmed when stress is extreme.

As noted in the introduction, one original rationale for including A2 in DSM-IV was that expansion of criterion A1 in DSM-IV might lead to an inappropriate broadening of the definition of PTSD unless a new subjective A2 requirement was used to delimit the range of qualifying traumas (2,4,5). As it turns out, the WMH results show that A2 did not have this desired effect. A somewhat different version of this argument is that the definition of a traumatic experience needs to include a short-term subjective component to focus clinical attention shortly after the time of trauma exposure on the subset of PTEs most likely to lead to the other PTSD criteria. This view might make good sense when it comes to predicting subsequent onset of PTSD from information about short-term emotional reactions to PTEs because conditional risk of PTSD is low in the absence of A2. Longitudinal studies have confirmed this predictive association (31,32). Targeting preventive interventions consequently might make use of such information (33).

This possible value of the assessment of A2 near the time of trauma exposure as a predictor of future PTSD should not be confused with the role of A2 as a requirement for a diagnosis of PTSD. The distinction is familiar in other areas of medicine. For example, detection of hypertension is clinically useful as a predictor of heart attack and usually triggers initiation of preventive interventions. It makes no sense, though, to require a history of hypertension for a diagnosis of myocardial infarction when a person has a heart attack. Similarly with PTSD, when a patient presents for treatment of reactions to a PTE that include DSM-IV criteria B through F, it would seem perverse to use the patient's retrospective report of not experiencing A2 at the time of the PTE exposure to rule out a diagnosis of PTSD.

Another argument for retaining A2 is that information about A2 can be valuable in making a differential diagnosis between PTSD and other syndromes, such as comorbid reactive depression with phobia, when a patient presents with a mixture of depression and anxiety related to a PTE. The question of whether to make a diagnosis of PTSD if the patient's current symptoms meet PTSD criteria B through F even if the patient fails to recall A2 emotional reactions at the time of the trauma then has to be considered. The data presented here are unable to address this issue. Research to investigate this issue would be an important addition to other studies that have been called for to refine our understanding of differential treatment response in PTSD (34), especially if a differential treatment response could be found based on retrospective assessment of A2.

These findings should be interpreted in the context of several important limitations. First, WMH interviews were conducted by lay interviewers rather than clinicians. This is an especially important issue for the evaluation of retrospective reports of complex emotional responses like those involved in A2. Second, the clinical validation of PTSD was restricted to the United States. Concordance of the CIDI diagnoses with clinical diagnoses may not be as good in some other countries. Third, PTSD was assessed in a lifetime framework, which introduces the possibility of recall bias. Fourth, results may not generalize to PTEs that

were not considered in the CIDI assessment or that were so rarely reported that few such PTEs were represented in our randomly selected series.

The third limitation is especially important in light of the high proportion of respondents with other requirements for a diagnosis of PTSD who retrospectively reported that they met A2 at the time of PTE exposure (98.6%). This proportion exceeds the comparable proportion found in the one prospective general population study that examined this issue (89%) (26), raising the possibility that A2 is recalled as being more prevalent than it actually was. If this is the case, then the small proportion of cases of PTSD estimated to be excluded from a diagnosis by the A2 requirement may be underestimated. This possibility has to be considered in the context of the fact that some investigators have called for A2 to be broadened beyond the focus on fear, helplessness, and horror to include other strong emotions sometimes associated with PTSD, including anger, shame, grief, and extreme emotional blunting (i.e., shock, dissociation) (26,35,36). If A2 is broadened in these ways, then it is easy to imagine that virtually 100% of the people who meet other requirements for PTSD would also meet the broadened definition of A2. This would make A2 completely redundant with the other criteria, which would argue even more forcefully than the results reported here that A2 should be considered a risk factor rather than a diagnostic criterion for PTSD.

This report is carried out in conjunction with the World Health Organization (WHO) World Mental Health Survey Initiative. We thank the World Mental Health staff for assistance with instrumentation, field work, and data analysis. These activities were supported by the United States National Institute of Mental Health (R01MH070884), the John D. and Catherine T. MacArthur Foundation, the Pfizer Foundation, the US Public Health Service (R13-MH066849, R01-MH069864, and R01 DA016558), the Fogarty International Center (R03-TW006481), the Pan American Health Organization, the Eli Lilly & Company Foundation, Ortho-McNeil Pharmaceutical Inc., GlaxoSmithKline, Bristol-Myers Squibb, and Shire. A complete list of World Mental Health publications can be found at <http://www.hcp.med.harvard.edu/umb/>. The São Paulo Megacity Mental Health Survey is supported by the State of São Paulo Research Foundation Thematic Project Grant 03/00204-3. The Bulgarian Epidemiological Study of Common Mental Disorders is supported by the Ministry of Health and the National Center for Public Health Protection. The Chinese World Mental Health Survey Initiative is supported by the Pfizer Foundation. The Colombian National Study of Mental Health is supported by the Ministry of Social Protection. The European Study Of The Epidemiology Of Mental Disorders project is funded by the European Commission (Contracts QLG5-1999-01042; SANCO 2004123); the Piedmont Region (Italy); Fondo de Investigación Sanitaria, Instituto de Salud Carlos III, Spain (FIS 00/0028); Ministerio de Ciencia y Tecnología, Spain (SAF 2000-158-CE); Departament de Salut, Generalitat de Catalunya, Spain, Instituto de Salud Carlos III (Centro de Investigación Biomédica en Red CB06/02/0046, Redes Temáticas de Investigación Cooperativa en Salud RD06/0011 Red de Enfermedades Mentales), and other local agencies and by an unrestricted educational grant from GlaxoSmithKline. The World Mental Health India was funded by WHO (India) and helped by Dr. R. Chandrasekaran, Jawaharlal Institute of Postgraduate Medical Education & Research. The Israel National Health Survey is funded by the Ministry of Health with support from the Israel National Institute for Health Policy and Health Services Research

and the National Insurance Institute of Israel. The World Mental Health Japan Survey is supported by the Grant for Research on Psychiatric and Neurological Diseases and Mental Health (H13-SHOGAI-023, H14-TOKUBETSU-026, H16-KOKORO-013) from the Japan Ministry of Health, Labour and Welfare. The Lebanese National Mental Health Survey LEBANON is supported by the Lebanese Ministry of Public Health, the WHO (Lebanon), Fogarty International, anonymous private donations to Institute for Development, Research, Advocacy & Applied Care, Lebanon, and unrestricted grants from Janssen Cilag, Eli Lilly, GlaxoSmithKline, Roche, and Novartis. The Mexican National Comorbidity Survey is supported by The National Institute of Psychiatry Ramon de la Fuente (Instituto Nacional de Psiquiatria Ramon de la Fuente Muniz Direccion de Investigaciones Epidemiologicas y Sociales 4280) and by the National Council on Science and Technology (CONACyT-G30544-H), with supplemental support from the PanAmerican Health Organization. Te Rau Hinengaro: The New Zealand Mental Health Survey is supported by the New Zealand Ministry of Health, Alcohol Advisory Council, and the Health Research Council. The Nigerian Survey of Mental Health and Wellbeing is supported by the WHO (Geneva), the WHO (Nigeria), and the Federal Ministry of Health, Abuja, Nigeria. The Romania World Mental Health study projects "Policies in Mental Health Area" and "National Study Regarding Mental Health and Services Use" were carried out by National School of Public Health & Health Services Management (former National Institute for Research & Development in Health), with technical support of Metro Media Transilvania, the National Institute of Statistics-National Centre for Training in Statistics, Statistics contract 70, Cheyenne Services Societate cu Responsabilitate Limitata, and Statistics Netherlands and were funded by Ministry of Public Health (former Ministry of Health) with supplemental support of Eli Lilly, Romania Societate cu Responsabilitate Limitata. The South Africa Stress and Health Study is supported by the US National Institute of Mental Health (R01-MH059575) and National Institute of Drug Abuse with supplemental funding from the South African Department of Health and the University of Michigan. The Ukraine Comorbid Mental Disorders during Periods of Social Disruption study is funded by the US National Institute of Mental Health (RO 1-MH61905). The US National Comorbidity Survey Replication is supported by the National Institute of Mental Health (U01-MH60220) with supplemental support from the National Institute of Drug Abuse, the Substance Abuse and Mental Health Services Administration, the Robert Wood Johnson Foundation (Grant 044708), and the John W. Alden Trust.

We thank Ms. Yasmine Fayyad for working on the literature review.

Dr. Kessler has been a consultant for GlaxoSmithKline Inc., Kaiser Permanente, Pfizer Inc., Sanofi-Aventis, Shire Pharmaceuticals, and Wyeth-Ayerst; has served on advisory boards for Eli Lilly & Company and Wyeth-Ayerst; and has had research support for his epidemiological studies from Bristol-Myers Squibb, Eli Lilly & Company, GlaxoSmithKline, Johnson & Johnson Pharmaceuticals, Ortho-McNeil Pharmaceuticals Inc., Pfizer Inc., and Sanofi-Aventis. Dr. Stein has received research grants and/or consultancy honoraria from AstraZeneca, Eli-Lilly, GlaxoSmithKline, Johnson & Johnson, Lundbeck, Orion, Pfizer, Pharmacia, Roche, Servier, Solway, Sumitomo, Tivvah, and Wyeth. The remaining authors report no biomedical financial interests or potential conflicts of interest.

1. Kilpatrick DG, Resnick HS, Freedy JR, Pelcovitz D, Resick PA, Roth S, *et al.* (1998): Posttraumatic stress disorder field trial: Evaluation of the PTSD construct—criteria A through E. In: Widiger TA, Frances AJ, Pincus HA, Ross R, First MB, Davis W, *et al.*, editors. *DSM-IV Sourcebook*, vol. 4. pp. 803–844. Washington, DC: American Psychiatric Publishing.
2. Weathers FW, Keane TM (2007): The criterion A problem revisited: Controversies and challenges in defining and measuring psychological trauma. *J Trauma Stress* 20:107–121.
3. American Psychiatric Association (1994): *Diagnostic and Statistical Manual of Mental Disorders, 4th ed.* Washington, DC: American Psychiatric Association.
4. Creamer M, McFarlane AC, Burgess P (2005): Psychopathology following trauma: The role of subjective experience. *J Affect Disord* 86:175–182.
5. McNally RJ (2004): Conceptual problems with the DSM-IV criteria for posttraumatic stress disorder. In: Rosen GM, editor. *Posttraumatic Stress Disorder: Issues and Controversies*. New York: Wiley, 1–14.
6. Breslau N, Kessler RC (2001): The stressor criterion in DSM-IV posttraumatic stress disorder: An empirical investigation. *Biol Psychiatry* 50:699–704.
7. Schnurr PP, Spiro A III, Vielhauer MJ, Findler MN, Hamblen JL (2002): Trauma in the lives of older men: Findings from the Normative Aging Study. *J Clin Geropsychol* 8:175–187.
8. O'Donnell ML, Creamer M, McFarlane AC, Silove D, Bryant RA (2010): Should A2 be a diagnostic requirement for posttraumatic stress disorder in DSM-V? *Psychiatry Res* 176:257–260.
9. Andrews G, Slade T, Sunderland M, Anderson T (2007): Issues for DSM-V: Simplifying DSM-IV to enhance utility: The case of major depressive disorder. *Am J Psychiatry* 164:1784–1785.
10. Zimmerman M, Chelminski I, McGlinchey JB, Young D (2006): Diagnosing major depressive disorder. X: Can the utility of the DSM-IV symptom criteria be improved? *J Nerv Ment Dis* 194:893–897.
11. Friedman MJ, Karam EG (2009): Posttraumatic stress disorder: Looking toward DSM-V and ICD-11. In: Andrews G, Charney DS, Sirovatka P, Regier DA, editors. *Stress-Induced and Fear Circuitry Disorders: Refining the Research Agenda for DSM-V*. Washington, DC: American Psychiatric Association Press, 3–24.
12. Kessler RC, Üstün TB (2008): The World Health Organization Composite International Diagnostic Interview. In: Kessler RC, Üstün TB, editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 58–90.
13. Heeringa SG, Wells JE, Hubbard F, Mneimneh Z, Chiu WT, Sampson N (2008): Sample designs and sampling procedures. In: Kessler RC, Üstün TB, editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 14–32.
14. Pennell B-E, Mneimneh Z, Bowers A, Chardoul S, Wells JE, Viana MC, *et al.* (2008): Implementation of the World Mental Health Surveys. In: Kessler RC, Üstün TB, editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 33–57.
15. Kessler RC, Üstün TB (2004): The World Mental Health (WMH) survey initiative version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int J Methods Psychiatr Res* 13:93–121.
16. Harkness J, Pennell B-E, Villar A, Gebler N, Aguilar-Gaxiola S, Bilgen I (2008): Translation procedures and translation assessment in the World Mental Health Survey Initiative. In: Kessler RC, Üstün TB, editors. *The WHO World Mental Health Surveys: Global Perspectives on the Epidemiology of Mental Disorders*. New York: Cambridge University Press, 91–113.
17. Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB (1995): Posttraumatic stress disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 52:1048–1060.
18. Breslau N, Kessler RC, Chilcoat HD, Schultz LR, Davis GC, Andreski P (1998): Trauma and posttraumatic stress disorder in the community: The 1996 Detroit Area Survey of Trauma. *Arch Gen Psychiatry* 55:626–632.
19. First MB, Spitzer RL, Gibbon M, Williams JBW (2002): *Structured Clinical Interview for DSM-IV Axis I Disorders, Research Version, Non-Patient Edition (SCID-I/NP)*. New York: Biometrics Research, New York State Psychiatric Institute.
20. Kessler RC, Abelson J, Demler O, Escobar JI, Gibbon M, Guyer ME, *et al.* (2004): Clinical calibration of DSM-IV diagnoses in the World Mental Health (WMH) version of the World Health Organization (WHO) composite International diagnostic interview (WMH-CIDI). *Int J Methods Psychiatr Res* 13:122–139.
21. Haro JM, Arbabzadeh-Bouchez S, Brugha TS, de Girolamo G, Guyer ME, Jin R, *et al.* (2006): Concordance of the Composite International Diagnostic Interview, Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO World Mental Health surveys. *Int J Methods Psychiatr Res* 15:167–180.
22. Hosmer DW, Lemeshow S (2004): *Applied Logistic Regression, 2nd ed.* New York: Wiley.
23. Efron B (1988): Logistic regression, survival analysis, and the Kaplan-Meier curve. *J Am Stat Assoc* 83:414–425.
24. Wolter KM (1985): *Introduction to Variance Estimation*. New York: Springer-Verlag.
25. Research Triangle Institute (2002): *SUDAAN: Professional Software for Survey Data Analysis, Version 8.0.1*. Research Triangle Park, NC: Research Triangle Institute.
26. Brewin CR, Andrews B, Rose S (2000): Fear, helplessness, and horror in posttraumatic stress disorder: Investigating DSM-IV criterion A2 in victims of violent crime. *J Trauma Stress* 13:499–509.
27. Janoff-Bulman R, Frantz CM (1996): The loss of illusions: The potent legacy of trauma. *J Pers Interpers Loss* 1:133–150.
28. Wortman CB, Silver RC (1989): The myths of coping with loss. *J Consult Clin Psychol* 57:349–357.
29. Adler AB, Wright KM, Bliese PD, Eckford R, Hoge CW (2008): A2 diagnostic criterion for combat-related posttraumatic stress disorder. *J Trauma Stress* 21:301–308.
30. Williams NL, Reardon JM, Murray KT, Cole TM (2005): Anxiety disorders: A developmental vulnerability-stress perspective. In: Hankin BL, Abela JRZ, editors. *Development of Psychopathology: A Vulnerability-Stress Perspective*. Thousand Oaks, CA: Sage, 289–327.
31. Ehrling T, Ehlers A, Cleare AJ, Glucksmann E (2008): Do acute psychological and psychobiological responses to trauma predict subsequent symptom severities of PTSD and depression? *Psychiatry Res* 161:67–75.
32. Zatzick DF, Kang SM, Muller HG, Russo JE, Rivara FP, Katon W, *et al.* (2002): Predicting posttraumatic distress in hospitalized trauma survivors with acute injuries. *Am J Psychiatry* 159:941–946.
33. Rauch SA, Hembree EA, Foa EB (2001): Acute psychosocial preventive interventions for posttraumatic stress disorder. *Adv Mind Body Med* 17: 187–191.
34. Stein DJ, Cloitre M, Nemeroff CB, Nutt DJ, Seedat S, Shalev AY, *et al.* (2009): Cape Town consensus on posttraumatic stress disorder. *CNS Spectr* 14:52–58.
35. Brunet A, Weiss DS, Metzler TJ, Best SR, Neylan TC, Rogers C, *et al.* (2001): The Peritraumatic Distress Inventory: A proposed measure of PTSD criterion A2. *Am J Psychiatry* 158:1480–1485.
36. Roemer L, Litz BT, Orsillo SM, Ehlich PJ, Friedman MJ (1998): Increases in retrospective accounts of war-zone exposure over time: The role of PTSD symptom severity. *J Trauma Stress* 11:597–605.